arXiv:math/0506404v2 [math.PR] 19 Dec 2006

Oriented Percolation in One–Dimensional $\beta/|x-y|^2$ Random Cluster Model at $\beta > 1$

D. H. U. Marchetti[*], V. Sidoravicius[†] and M. E. Vares[‡]

Abstract

We consider the Fortuin-Kasteleyn random cluster model on $\mathbb{Z}$, with edge occupation probabilities

$$p_{\{x,y\}} = \begin{cases} p & \text{if } |x-y| = 1, \\ 1 - \exp\left\{-\beta/\left|x-y\right|^2\right\} & \text{otherwise,} \end{cases}$$

and weighting factor $\kappa \geq 1$. We prove that oriented percolation occurs when $\beta > 1$ and $\kappa \geq 1$ provided p is chosen sufficiently close to 1. In the particular case of the product measure ($\kappa = 1$), this answers a question posed in [NS]. The proof is based on multi-scale analysis.

1 Introduction

It is well known that $1/r^2$ gives the "critical" falloff for percolation in one-dimensional long range independent edge percolation models. Moreover, for the one dimensional Fortuin–Kasteleyn (FK) random cluster model with weighting factor $\kappa \geq 1$ and edge occupation probabilities of the form $p_{\{x,y\}} = f(|x-y|)$, with $\beta := \lim_{x\to\infty} x^2 f(x) > 0$, it is known that for fixed $f(j) < 1, j \geq 2$ and varying $p = f(1)$, the value $\beta^* = 1$ is critical in the sense that for $\beta \leq 1$ percolation cannot occur unless $p = 1$ (see [AN]), while for $\beta > 1$ there is percolation provided p is sufficiently close to one (see [IN] and [M]). These results are important in the description of the phase transition diagram for the one-dimensional long range Ising models studied earlier by Fröhlich and Spencer in [FS1] and for the corresponding Potts models ([ACCN],[IN],[M]), as these spin systems can be constructed by a random coloring of the clusters in the FK model with $\kappa = 2$, or $\kappa > 2$ integer, respectively. For the particular case of independent edge percolation models ($\kappa = 1$) earlier results were obtained in [NS], where it was proven that $\beta^* \leq 1$ in this case, and that *oriented percolation* occurs when $\lim_{x\to\infty} x^s f(x) > 0$ for some $1 < s < 2$. The question whether oriented percolation occurs in the boundary case $s = 2$ remained unanswered. Theorem 1.1 below gives an affirmative answer applicable to the FK model: the result is stated for the particular example of edge probabilities

---

[*] Instituto de Física, Universidade de São Paulo, Caixa Postal 66318, 05315 São Paulo, SP, Brasil. E-mail: *marchett@if.usp.br*

[†] Instituto de Matemática Pura e Aplicada (IMPA), Estrada Dona Castorina 110, Jardim Botânico, 22460-320 Rio de Janeiro, RJ, Brasil. E-mail: *vladas@impa.br*

[‡] Centro Brasileiro de Pesquisas Físicas (CBPF), Rua Dr. Xavier Sigaud 150, 22290-180 Rio de Janeiro, RJ, Brasil. E-mail: *eulalia@cbpf.br*

in (1.1) below, and oriented percolation is shown when $\beta > 1$ and $p < 1$ is sufficiently close to one. The proofs can be easily adapted to include any $f(\cdot)$ satisfying $\lim_{x\to\infty} x^2 f(x) > 1$. In this sense $\beta^* = 1$ remains critical also for oriented percolation.

Preliminaries. Consider the infinite complete graph with set of vertices $\mathcal{V} = \mathbb{Z}$ and set of edges $\mathbb{E} = \{\{x, y\},\ x \neq y,\ x, y \in \mathbb{Z}\}$, and let $\Omega = \{0,1\}^{\mathbb{E}}$. One-dimensional long-range FK random cluster models with weighting parameter $\kappa > 0$ are probability measures on $\sigma(\Omega)$, the product σ–algebra on Ω. To define them, let us first fix ν the Bernoulli product measure on Ω, with $\nu(\omega_{\{x,y\}} = 1) = p_{\{x,y\}}$ given by

$$p_{\{x,y\}} = \begin{cases} p & \text{if } |x-y| = 1, \\ 1 - \exp\left\{-\dfrac{\beta}{|x-y|^2}\right\} & \text{otherwise,} \end{cases} \tag{1.1}$$

where $0 < p < 1$ and $\beta > 0$ are fixed parameters.

Notation. We write $q_{\{x,y\}} = 1 - p_{\{x,y\}}$; for $e = \{x, y\}$ we will write p_e instead of $p_{\{x,y\}}$, and say that e "is open" if $\omega_e = 1$. The length of an edge $e = \{x, y\}$ is $|x - y|$.

Finite volume FK measure. Given $I \subset \mathbb{Z}$, consider $\mathbb{E}(I) = \{\{x, y\} \in \mathbb{E}\colon x, y \in I\}$, $\Omega_I = \{0,1\}^{\mathbb{E}(I)}$, and $\bar\Omega_I = \{0,1\}^{\mathbb{E}\setminus\mathbb{E}(I^c)}$, where $I^c = \mathbb{Z}\setminus I$. Assume that $|I| < \infty$, and fix an "external" configuration $\tilde\omega \in \{0,1\}^{\mathbb{E}(I^c)}$. The finite volume FK-measure with external conditions $\tilde\omega$ is the probability measure $\mu^{\tilde\omega}_{\kappa,I}$ on $\bar\Omega_I$, given by

$$\mu^{\tilde\omega}_{\kappa,I}(A) = \frac{\displaystyle\int_A \kappa^{\mathcal{C}_I(\omega\circ\tilde\omega)}\ \bar\nu_I\,(d\omega)}{\displaystyle\int_{\bar\Omega_I} \kappa^{\mathcal{C}_I(\omega\circ\tilde\omega)}\ \bar\nu_I\,(d\omega)}\ , \tag{1.2}$$

where $\bar\nu_I$ is the projection of ν on $\bar\Omega_I$, the configuration $\omega\circ\tilde\omega$ is given by $(\omega\circ\tilde\omega)_e = \omega_e$ if $e \in \mathbb{E}\setminus\ \mathbb{E}(I^c)$ and $(\omega \circ \tilde\omega)_e = \tilde\omega_e$ if $e \in\ \mathbb{E}(I^c)$, and for any configuration $\omega \in \Omega$, $\mathcal{C}_I(\omega)$ is the number of disjoint connected components intersecting I in the graph determined by this configuration (i.e. the graph whose edges coincide with those e such that $\omega_e = 1$).

Of course we may think of $\mu^{\tilde\omega}_{\kappa,I}$ as a probability measure on $\Omega^{\tilde\omega}_I = \{\omega \in \Omega : \omega_e = \tilde\omega_e, \text{if } e \in \mathbb{E}(I^c)\}$.

Several delicate issues are present in the passage to the infinite volume limit, specially when $0 < \kappa < 1$. (See [G1], [G2].)

Throughout the paper we will assume that $\kappa \geq 1$. Two external configurations $\tilde\omega$ play a particular role when considering the infinite volume limit:

$\tilde\omega_e = 0$ for all $e \in \mathbb{E}(I^c)$, and the corresponding measure $\mu^{\tilde\omega}_{\kappa,I} \equiv \mu^{f}_{\kappa,I}$ is called the "free" measure;

and

$\tilde\omega_e = 1$ for all $e \in \mathbb{E}(I^c)$, and the corresponding measure $\mu^{\tilde\omega}_{\kappa,I} \equiv \mu^{w}_{\kappa,I}$ is called the "wired" measure.

The infinite volume limit. On Ω we consider the usual partial order: $\omega \leq \omega'$ if $\omega_e \leq \omega'_e$ for each $e \in \mathbb{E}$. By the FKG inequality (see [F], [ACCN]), one has

$$\mu^{f}_{\kappa,I}(g) \leq \mu^{f}_{\kappa,I'}(g) \leq \mu^{\tilde\omega}_{\kappa,I'}(g) \leq \mu^{w}_{\kappa,I'}(g) \leq \mu^{w}_{\kappa,I}(g)$$

for any finite intervals $I \subset I' \subset \mathbb{Z}$, any external configuration $\tilde{\omega}$ and any non-decreasing continuous function $g : \Omega \to \mathbb{R}$. Thus, as $I \nearrow \mathbb{Z}$ the limit measures μ_κ^f and μ_κ^w exist. Moreover, $\mu_\kappa^f \le \mu_\kappa^w$ in FKG sense[1], and $\mu_\kappa^f = \mu_\kappa^w$ if and only if there is a unique accumulation point of the set $\mu_{\kappa,I}^{\tilde{\omega}}$ as $I \nearrow \mathbb{Z}$. If $\kappa = 1$, trivially $\mu_\kappa^f = \mu_\kappa^w = \nu$. Since $p_{\{x,y\}} \equiv f(|x-y|)$ the measures μ_κ^f and μ_κ^w are translation invariant; both these measures are ergodic. For a more general discussion on the construction of random cluster measures see e.g. [G1].

Fix $\omega \in \Omega$. An alternating sequence of vertices and edges $x = x_1, e_1, x_2, \dots, x_{n-1}, e_{n-1}, x_n = y$, $n \ge 1$, is called a path connecting x to y, and we say that the path is open if $\omega_{e_i} \equiv \omega_{\{x_i,x_{i+1}\}} = 1$, $1 \le i \le n-1$. We say that $C \subset \mathbb{Z}$ is connected if for any two distinct vertices x, y in C there exists an open path π connecting them. A maximal connected set is called an *open cluster*, and $C_x(\omega)$ denotes the open cluster containing $x \in \mathbb{Z}$ (we write $C_x(\omega) = \{x\}$ if $\omega_{\{x,y\}} = 0$, for all $y \in \mathbb{Z} \setminus \{x\}$). A path $\pi = (x_1, \dots, x_n)$ connecting x to y, $x < y$, is called *oriented* if $x_1 = x < x_2 < \cdots < x_{n-1} < x_n = y$, and we write $x \rightsquigarrow y$ when there is an open oriented path connecting x to y. Analogously we define $C_x^+ = \{y \colon x \rightsquigarrow y\}$, and the event

$$[x \rightsquigarrow \infty] = [|C_x^+| = \infty].$$

For $L \ge 1$ integer, and use $\mu_{\kappa,L}^{\tilde{\omega}}$ as a shorthand for $\mu_{\kappa,[-L,L]}^{\tilde{\omega}}$.

We are ready to state our main result.

Theorem 1.1 *For any* $\beta > 1$, $\kappa \ge 1$, *there exist* $0 < p_0 < 1$ *such that, if* $p > p_0$, *then*

$$\mu_\kappa^f(0 \not\rightsquigarrow \infty) \le \epsilon, \tag{1.3}$$

with $\epsilon = \epsilon(p) \searrow 0$ *as* $p \nearrow 1$.

Remarks. **1**. Since $\mu_\kappa^f \le \mu_\kappa^w$, the previous statement holds as well for μ_κ^w.

2. The proof of this theorem can be adapted to establish an analogue of Theorem 1.2 of [M] on the power law decay of truncated connectivity function, $\tau'(x) := \mu_\kappa^w(0 \rightsquigarrow x, 0 \not\rightsquigarrow \infty)$ (see [IN] and Remarks after Theorem 1.2 in [M] for the significance of this quantity in the non-oriented case). To get the right decay is still an open problem in the oriented case.

Some related problems. The type of questions treated here has various sources of interest and we mention only a couple of them, which have to do with our own motivations. Consider the following physical problem: take the one-dimensional Ising model with pair interactions, the couplings decaying as the square of the distance between vertices, at inverse temperature $\beta > 1$; this is the model studied by Fröhlich and Spencer ([FS1]), for which a phase transition was established. Take now the finite box $[-L, L]$ and assume the Dobrushin boundary conditions, i.e. all spins in $(-\infty, -L]$ will be taken as $+1$, and all spins in $[L, +\infty)$ will be taken as -1. What can we say about the behaviour of this model when $L \to \infty$? Is there any sort of well defined *interface*? This can be rephrased in a more general form in the language of the FK representation as a question about the behavior of connected components of each boundary in the corresponding dependent percolation model, conditioned not to touch each other. (Recall that by a random coloring of the clusters, the FK model gives origin to a spin system which interpolates the independent percolation model ($\kappa = 1$), the Ising model ($\kappa = 2$) and the q–states Potts model ($\kappa = q > 2$, integer) at inverse temperature β and interaction $J_{\{x,y\}} = \beta^{-1}\log(\frac{1}{1-p_{\{x,y\}}})$. For details see [FK, ACCN]). Though we still do not fully understand this problem, our results shed some light on it and hint that typically the connected cluster of one of the sides $(-\infty, -L]$ or $[L, +\infty)$, (w.p. close to 1/2, each) would take over most of the "middle" space. This is an unsolved problem. In [CMR], the authors obtain a more precise description for very low temperatures, using cluster expansion techniques.

---

[1] That is $\mu \le \mu'$ if $\mu(g) \le \mu'(g)$ for any g continuous and increasing.

An interesting corollary of Theorem 1.1 is as follows. Consider the Ising model (with $\pm$–valued spins) with interaction $J_{\{x,y\}} = |x-y|^{-2}$ if $|x-y| \geq 2$ and $J_{\{x,x+1\}} = J$ at inverse temperature β. Let $m_L^{0,+}(\beta)$ denote the average spin at the origin, with "one-sided" $(+)$ boundary conditions in $[L,\infty)$. By the above mentioned FK representation, we have

Corollary 1.2 *Under the conditions of Theorem 1.1,*

$$m_L^{0,+}(\beta) \geq \mu^w_{2,L}(0 \rightsquigarrow \infty) \geq 1-\epsilon$$

holds for $\epsilon = \epsilon(J) \searrow 0$ *as* $J \nearrow \infty$ *uniformly in* L*. Consequently, there exists a phase transition when the thermodynamical limit is taken with free boundary conditions on the left side of the interval* $[-L, L]$*, and* $+$ *boundary conditions on the right side.*

It is also interesting to compare the result on oriented percolation and the previous corollary with the somehow similar question on the multiplicity of Gibbs states for Markov chains with infinite connections, where orientation appears naturally through the time direction. Recently Johansson and Öberg [JO] showed that if g is a regular specification and

$$\mathrm{var}_k(g) = \sup\{\|g(\sigma) - g(\sigma')\|_1 \colon \sigma_i = \sigma'_i, i = 1, \ldots, k\},$$

then g admits a unique Gibbs measure whenever the sequence $\{\mathrm{var}_k(g)\}_{k=1}^{+\infty}$ is in ℓ^2. This tells, in particular, that there are no multiple limiting measures for chains with connections decaying as r^{-2}, as in Example 1 in [JO]. This contrasts with the two-sided Ising models and, as our Theorem says, with percolation models. The understanding of Markov chains with infinite connections in the non-uniqueness regime is still very poor, and it is known as a notoriously difficult problem. We have strong evidence (see [BHS]) that multi-scale analysis techniques analogous to those developed in this work could be turned into a robust tool to study this question.

Heuristics of the proof. The proof relies on Fröhlich-Spencer multi-scale analysis ideas ([FS], [FS1]), and we use the version developed in [KMP] and [M]. In the next few paragraphs we outline the scheme of the proof, and comment on some key ideas, avoiding most of consuming technical points. Our goal here is only to give a very schematic and approximate picture, postponing precise formulations (which tend to be quite involved) to later in the text.

The goal. We look for an event of positive probability, whose occurrence implies not only the existence of an infinite open component, but also guarantees the presence of an oriented infinite open path. Essentially, we will construct such an event, and show that it has positive probability. Our key estimate will be: if $\beta > 1$, we can find $\delta > 0$, $\delta' < 1$ and p is sufficiently close to 1, so that

$$\nu(\exists \,\text{open path}\; \pi = (x_1, e_1, \ldots, x_n) \colon x_1 \leq -L + L^{\delta'},\, x_n \geq L - L^{\delta'}, 0 < x_i - x_{i-1} \leq L^{\delta'},\, \forall i\,) \geq 1 - L^{-\delta}$$

for $L = l_k$ as defined below $(k \geq 1)$, l_1 being sufficiently large, and where ν stands for the product measure defined before. Further control for the dependent model leading to the result for μ^f_κ will come along the proof. We will have little control on how close to one p has to be (or, on how large l_1 we need).

Scales. We choose super-exponentially fast growing scales. Given $1 < \alpha < 2$, $l_0 = 1$ and l_1 an integer sufficiently large, let

$$l_k = \lfloor l_{k-1}^{\alpha-1} \rfloor l_{k-1}, \qquad k = 2, 3, \dots, \tag{1.4}$$

where as usual $\lfloor z \rfloor = \max\{n \in \mathbb{N} \colon n \leq z\}$. We will use the so–called dynamical blocking argument, in that the size and location of blocks[2] will be defined along the procedure and will depend on the configuration. Still, the length of each block $I^{(k)}$ of the k–th level (called k–block) will be of order l_k. More precisely, we shall see that $l_k - 2l_k^{\alpha'/\alpha} - 6l_{k-1} \leq |I^{(k)}| \leq 3l_k + 6l_{k-1}$, for suitable $1 < \alpha' < \alpha$. (In particular, $l_{k-1} \ll l_k^{\alpha'/\alpha} \ll l_k$, if $k \geq 2$, and l_1 is large.)

Defected and good blocks. Further we will use the following recursive definition of a "defected" block. Fix $1 < \alpha' < \alpha$ to be specified later.

1) We say that the 0–block $[i, i+1]$ is *defected* if the corresponding nearest neighbor edge $\{i, i+1\}$ is closed; otherwise the 0–block is said to be *good* and the open nearest neighbor path from i to $i+1$ is called a 0–*pedestal*;

2) For $k \geq 1$, a k–block $I^{(k)} = [s, s']$ is *defected* if either it contains two or more defected $(k-1)$–blocks, or it contains only one defected $(k-1)$–block $[i, i']$ but there is no open edge $\{a, a'\}$ of length at most $l_k^{\alpha'/\alpha}$, with $a \leq i$, $i' \leq a', a \in \Upsilon$, $a' \in \Upsilon'$, for some $(k-1)$–pedestals Υ, Υ' contained in $I^{(k)}$. Otherwise $I^{(k)}$ is called *good.*

Thus, if a k–block $[s, s']$ is good, then it contains an oriented open path going from s to s': in the case it has no defected $(k-1)$–blocks, this path can be obtained by concatenating $(k-1)$–pedestals of the good $(k-1)$–blocks which constitute the given k–block; if it has a (single) defected $(k-1)$–block, and using the above notation, a similar concatenation yields an oriented open path going from s to a, which is followed by an open edge $\{a, a'\}$, and then followed by another concatenation of $(k-1)$–pedestals of good $(k-1)$–blocks, from a' to s'. In both cases, such path from s to s' will be called *k–pedestal*, and denoted by Υ. The part of the cluster between a and a' is again disregarded in the future construction since we have little control on oriented connectivity in this segment. The condition $a' - a \leq l_k^{\alpha'/\alpha}$ will be crucial to guarantee that pedestals are quite dense sets (within the corresponding good blocks), used to push the construction to higher levels. Some care is needed when treating defects close to the boundary, which we have disregarded here.

Strategy. Being "defected" doesn't necessarily imply that there is no oriented open path connecting the endpoints of the block. Nevertheless, in order to avoid substantial technical difficulties, we will follow two rules that simplify our construction:

a) once a block is defected, we will assume the worst possible situation, namely it will be considered as if all edges within this block are closed.

b) once we have at least two defected $(k-1)$–blocks within a k–block I, we will not try to find connections within the k–block to fix its connectivity, but rather will "push the problem to the next level", and try to "jump over" this troubled block I by a longer edge of length at most $l_{k+1}^{\alpha'/\alpha}$, which starts at the pedestal of some good k–block to the left of I, and ends similarly on the right of I.

---

[2] Successive blocks share an end-vertex.

Estimates. The scale l_1 will be taken large enough, as to be determined later. Assume that $\kappa \geq 1$ is fixed. We begin by choosing p:

$$p \geq 1 - \kappa^{-2} \frac{(\ln 2)^5}{128} \, l_1^{-\delta-1} \tag{1.5}$$

for some $\delta > 0$ to be chosen later (see (2.19) below). This particular dependence on κ will allow us to extend the estimate from the product measure ν to the random cluster measure μ_κ^f.

For $k \geq 2$, let $I^{(k)}$ be a k–block of length[3] l_k, which consists of $N_k = l_k / l_{k-1} = \lfloor l_{k-1}^{\alpha-1} \rfloor$ blocks of level $(k-1)$, of length l_{k-1}, and written as $\{I_j^{(k-1)}\}_{j=1}^{N_k}$. Assume that we have the following estimate

$$\nu(I_j^{(k-1)} \text{ is defected}) \leq l_{k-1}^{-\delta}, \quad 1 \leq j \leq N_k.$$

Under the above assumptions, and if δ is chosen to satisfy (2.19), we see that

$$\nu\Big(\exists\, 1 \leq i < j \leq N_k \colon I_i^{(k-1)}, I_j^{(k-1)} \text{ are both defected}\Big) \leq l_k^{-\delta}.$$

When the defected $I_i^{(k-1)}$ is unique, we assume for the moment that it stays at distance larger than $l_k^{\alpha'/\alpha}$ from the boundary of $I^{(k)}$. (Otherwise a sequence of local adjustments of blocks will be needed, as we shall see in Sect. 2. The left- and right-most extremal blocks in our volume are treated differently.) In this case let a and a' be the end-vertices of the unique defected block $I_i^{(k-1)}$. By our construction, there exists an oriented path starting from the left boundary of $I^{(k)}$ and ending at the vertex a and another open oriented path starting from vertex a' and going to the right boundary of $I^{(k)}$. Both these paths are obtained by concatenating pedestals of all good $(k-1)$–blocks on the left side of the defected block $I_i^{(k-1)}$ and, respectively, on the right side. We denote these new left and right pedestals by Υ and Υ', respectively. Given that $I^{(k)}$ has a unique defected $I_i^{(k-1)} = [a, a']$, and the pedestals Υ and Υ', one has the following upper bound for the conditional probability of not finding an open edge $\{x, y\}$ with $x \leq a, a' \leq y,\ x \in \Upsilon,\ y \in \Upsilon'$ and $y - x \leq l_k^{\alpha'/\alpha}$:

$$\prod_{\substack{x,y: x \leq a < a' \leq y, \\ y - x < l_k^{\alpha'/\alpha} \\ x \in \Upsilon,\, y \in \Upsilon'}} q_{\{x,y\}} = \exp\Big\{ - \sum_{\substack{x,y: x \leq a < a' \leq y, \\ y - x < l_k^{\alpha'/\alpha} \\ x \in \Upsilon,\, y \in \Upsilon'}} \frac{\beta}{|x-y|^2} \Big\} \leq l_{k-1}^{-\beta(1-\eta)(\alpha'-1)}, \tag{1.6}$$

where $\eta = \eta(\alpha, \alpha', l_1) > 0$, and can be taken arbitrarily small if $l_1 \to \infty$.

The precise statement and proof of the above estimate will be given in Lemma 2.1. It requires some work, and in order to obtain it for suitable $\eta = \eta(\alpha, \alpha', l_1) > 0$ which can be taken arbitrarily small if $l_1 \to \infty$ we will need to use certain geometric properties of pedestals Υ and Υ', which propagate inductively from each level into the next one. Namely, the pedestals are relatively dense sets (see (2.9) in Sect. 2) as the construction will show. Using the above estimate, writing

$$\big\{ I^{(k)} \text{ has a unique defected } (k-1)\text{–block } [a, a'] \text{ but remains defected } \big\} =$$

$$\big\{ I^{(k)} \text{ has unique defected } (k-1)\text{–block } [a, a'] \big\} \cap$$

$$\big\{ \text{ there is no open edge } \{x, y\} \text{ with } x \leq a, a' \leq y,\ x \in \Upsilon,\ y \in \Upsilon' \text{ and } y - x \leq l_k^{\alpha'/\alpha} \ \big\} ,$$

---

[3] This is not exact in general, but holds approximately.

and since these events depend on disjoint sets of edges, we easily get:

$$\nu\big(I^{(k)} \text{ has a unique defected } (k-1)\text{–block but remains defected}\big) \le l_{k-1}^{\alpha-1-\delta} l_{k-1}^{-\beta(1-\eta)(\alpha'-1)} \le l_k^{-\delta},$$

provided

$$\beta(1-\eta)(\alpha'-1) > (\delta+1)(\alpha-1). \tag{1.7}$$

Since $\beta > 1$ and $\eta = \eta(\alpha, \alpha', l_1)$ can be taken very small provided l_1 is large, it will suffice to suitably fix the parameters α and α' (α' close enough to α). This is done at the end of Sect. 2.

Difficulties. To carry on this scheme we have to go through several "unpleasant" and rather involved points. The use of a dynamical blocking argument, with the blocks of a given level depending not only on the size and location of lower level blocks, but also on their "status" (defected or good), requires a rather tight bookkeeping. This is expressed through what we call "itineraries".

The second difficulty is in the representation of the initial measure, which should be sensitive to the above mentioned dynamic renormalization. We opt to write it as a convex combination of a family of probability measures associated with the above mentioned itineraries.

Once this is achieved, all necessary estimates follow along the scheme of [FS1] and [KMP].

The plan of the paper goes as follows: in Section 2 we define the blocks and describe the dynamic renormalization procedure. In Section 3 we see how the product measure can be rewritten to take advantage of the previously described blocks in order to recursively estimate the probability of having a good k–block. Finally, in Section 4 we see how to apply the previous estimates to conclude Theorem 1.1.

Acknowledgments The authors thank C. M. Newman for suggesting this problem and A.-S. Sznitman for many useful discussions. We thank E. Presutti for pointing out the "interface question" and M. Cassandro and I. Merola for discussions on this topic. We also thank the referees for their careful reading and for their suggestions which helped us to improve the paper. This work was supported by Faperj grant E-26/151.905/2000, Fapesp grant 03/01366-7, CNPq - 477259/01-4, Fundacion Andes, and individual CNPq grants. We also thank various institutions CBPF, IMPA, IF-USP, ETH, UC-Berkeley and MSRI for financial support and hospitality during periods when this work was carried out.

2 Spatial blocks (Dynamic Renormalization)

Notation. For $L \in \mathbb{N}$, the construction will involve the configuration ω restricted to the set of edges with both end-vertices in $[-L, L]$. This allows us to prove the result for the measure μ_κ^f. We write Ω_L as a shorthand for $\Omega_{[-L,L]}$. Scales $\{l_k\}_{k\in\mathbb{N}}$ are defined in the following way: $l_0 = 1$, l_1 is an integer large enough to be chosen later (see (2.18) at the end of this Section), and for $1 < \alpha < 2$, we let

$$l_k = \lfloor l_{k-1}^{\alpha-1} \rfloor l_{k-1}, \quad k \ge 2. \tag{2.1}$$

Further we denote $x_j^{(k)} = j l_k,\ j \in \mathbb{Z}$.

For the proof of Theorem 1.1 we may assume that $L = l_M$, for some $M \in \mathbb{N}$.

Throughout the text $\mathbb{I}_A$ stands for the indicator function of an event A, i.e. $\mathbb{I}_A(\omega) = 1$ or 0 according to $\omega \in A$ or not.

Decomposition of events. Level 0. We set $I_i^{(0)} = [i, i+1]$. They are called 0–blocks, and for i such that $I_i^{(0)} \subset [-L, L]$ we define the events:

$$G(I_i^{(0)}) = \{\omega : \omega_{\{i,i+1\}} = 1\}, \qquad B(I_i^{(0)}) = \{\omega : \omega_{\{i,i+1\}} = 0\}.$$

$I_i^{(0)}$ is said to be defected when $B(I_i^{(0)})$ occurs; otherwise it is said to be a good 0–block.

Level 1. Consider the intervals $\widetilde{I}_j^{(1)} = [jl_1, (j+1)l_1]$ and for each j such that $\widetilde{I}_j^{(1)} \subset [-L, L]$ we define the following partition of Ω_L:

$$\begin{aligned}
G(\widetilde{I}_j^{(1)}) &= \bigcap_{i=jl_1}^{(j+1)l_1-1} G(I_i^{(0)}), \\
H_i(\widetilde{I}_j^{(1)}) &= B(I_i^{(0)}) \cap \bigcap_{\substack{s=jl_1 \\ s\neq i}}^{(j+1)l_1-1} G(I_s^{(0)}) \quad \text{for } i \in [jl_1, (j+1)l_1 - 1], \\
H(\widetilde{I}_j^{(1)}) &= \bigcup_{i=jl_1}^{(j+1)l_1-1} H_i(\widetilde{I}_j^{(1)}), \\
B(\widetilde{I}_j^{(1)}) &= \left(G(\widetilde{I}_j^{(1)}) \cup H(\widetilde{I}_j^{(1)})\right)^c, \qquad (2.2)
\end{aligned}$$

where G stands for *good*, H for *hopeful* and B for *bad*, and accordingly, $\widetilde{I}_j^{(1)}$ is said to be good (for given ω) if it contains no defected 0–blocks, "hopeful" if it contains only one defected 0–block, and is said to be "bad" otherwise. When $H_i(\widetilde{I}_j^{(1)})$ occurs, $I_i^{(0)}$ is called the defected 0-block in $\widetilde{I}_j^{(1)}$.

Adjustment. Given ω, we first consider the set of all j's such that $\omega \in H_{i_j}(\widetilde{I}_j^{(1)}) \subset H(\widetilde{I}_j^{(1)})$ and such that the index i_j of the (unique) defected block $I_{i_j}^{(0)} \subset \widetilde{I}_j^{(1)}$ verifies $jl_1 \le i_j \le jl_1 + \lfloor l_1^{\alpha'/\alpha} \rfloor - 1$ (resp. $(j+1)l_1 - \lfloor l_1^{\alpha'/\alpha} \rfloor \le i_j \le (j+1)l_1 - 1$).

If this set is empty in both cases, we set $I_j^{(1)} = \widetilde{I}_j^{(1)}$ for all j's, and say that $G(I_j^{(1)})$, $H(I_j^{(1)})$ or $B(I_j^{(1)})$ occurs if $G(\widetilde{I}_j^{(1)})$, $H(\widetilde{I}_j^{(1)})$ or $B(\widetilde{I}_j^{(1)})$ do occur.

If this set is not empty, we take arbitrarily one of such indices j; if $\widetilde{I}_j^{(1)}$ is not the interval which contains $-L$ (resp. L), to be treated in case 3) below, we check if $\widetilde{I}_{j-1}^{(1)}$ (resp. $\widetilde{I}_{j+1}^{(1)}$) has a defected 0–block in the sub-interval $[jl_1 - 2\lfloor l_1^{\alpha'/\alpha} \rfloor, jl_1]$ (resp. $[(j+1)l_1, (j+1)l_1 + 2\lfloor l_1^{\alpha'/\alpha} \rfloor - 1]$).

1) If yes, then we consider a new interval $I_{j-1}^{(1)} = \widetilde{I}_{j-1}^{(1)} \cup \widetilde{I}_j^{(1)}$ (resp. $I_j^{(1)} = \widetilde{I}_j^{(1)} \cup \widetilde{I}_{j+1}^{(1)}$) and say that the event $B(I_{j-1}^{(1)})$ (resp. $B(I_j^{(1)})$) occurs. (This is motivated by the fact that for the chosen ω the new interval will contain at least two defected 0–blocks.)

2) If not, then we consider two new intervals $I_{j-1}^{(1)} = [(j-1)l_1, jl_1 - \lfloor l_1^{\alpha'/\alpha} \rfloor]$ and $I_j^{(1)} = [jl_1 - \lfloor l_1^{\alpha'/\alpha} \rfloor, (j+1)l_1]$ (resp. $I_j^{(1)} = [jl_1, (j+1)l_1 + \lfloor l_1^{\alpha'/\alpha} \rfloor]$ and $I_{j+1}^{(1)} = [(j+1)l_1 + \lfloor l_1^{\alpha'/\alpha} \rfloor, (j+2)l_1]$) and

say that $G(I^{(1)}_{j-1})$, $H(I^{(1)}_{j-1})$ or $B(I^{(1)}_{j-1})$ occurs if the event $H(I^{(1)}_j)$ occurs and, resp. $G(\widetilde{I}^{(1)}_{j-1})$, $H(\widetilde{I}^{(1)}_{j-1})$ or $B(\widetilde{I}^{(1)}_{j-1})$, do occur (resp. $G(I^{(1)}_{j+1})$, $H(I^{(1)}_{j+1})$ or $B(I^{(1)}_{j+1})$ occurs if the event $H(I^{(1)}_j)$ occurs and, resp. $G(\widetilde{I}^{(1)}_{j+1})$, $H(\widetilde{I}^{(1)}_{j+1})$ or $B(\widetilde{I}^{(1)}_{j+1})$ do occur). In this case the adjustment moves the boundary "away" from the unique defected block in $I^{(1)}_j$, but doesn't change the number of the defected 0–blocks in the adjusted intervals.

3) If the interval $\widetilde{I}^{(1)}_j$ under consideration is the leftmost (resp. the rightmost) interval in $[-L, L]$, we set $I^{(1)}_j = \widetilde{I}^{(1)}_j$ and say that $G(I^{(1)}_j)$ occurs.

4) We set $I^{(1)}_j = \widetilde{I}^{(1)}_j$ if $\widetilde{I}^{(1)}_j$ was not involved in the previous adjustment, and say that $G(I^{(1)}_j)$, $H(I^{(1)}_j)$ or $B(I^{(1)}_j)$ occurs if accordingly $G(\widetilde{I}^{(1)}_j)$, $H(\widetilde{I}^{(1)}_j)$, or $B(\widetilde{I}^{(1)}_j)$ occur.

If after this step we are still left with intervals $I^{(1)}_j$ for which $H(I^{(1)}_j)$ occurs and its defected 0–block stays within distance $\lfloor l_1^{\alpha'/\alpha} \rfloor$ from the boundary of $I^{(1)}_j$, we repeat the above procedure, now applied to the intervals already adjusted in the previous step. After finitely many steps of such adjustment procedure there are left no intervals $I^{(1)}_j$ for which the event $H(I^{(1)}_j)$ occurs and its defected 0–block stays within distance $\lfloor l_1^{\alpha'/\alpha} \rfloor$ from the boundary, and the adjustment procedure is then stopped.

Remark. Notice that the adjustment procedure is well defined, i.e. the final partition does not depend on the order in which we do adjustments and in which order we pick the intervals that still need to be adjusted (in case if we have more than one). It also has *a locality* property, i.e. the final modification of each initial interval $\widetilde{I}^{(1)}_j$ depends on the values of the configuration in the nearest neighbor and, at most, in the next nearest neighbor intervals only.

Once the adjustment is completed, the obtained intervals are called 1-blocks, and re-numerated from left to right as $I^{(1)}_j$, $j = 1, \ldots$. Notice that $l_1 - 2\lfloor l_1^{\alpha'/\alpha} \rfloor \le |I^{(1)}_j| \le 3l_1$, and $\cup_j I^{(1)}_j = [-L, L]$.

In other words, values of ω on the nearest neighbor edges of $[-L, L]$ determine, through the above procedure, a division $I^{(1)}(\omega) \equiv \{I^{(1)}_j(\omega)\}_j$ of the interval $[-L, L]$ into 1–blocks, with the property that any two adjacent blocks are sharing their end-vertex. This is the final state of the "adjustment" procedure. Values of ω on the nearest neighbor edges in $[-L, L]$ also determine where the defected 0-blocks are located within each 1-block, and we denote by $D^{(1)}_j(\omega)$ the set of indices of the defected 0–blocks within $I^{(1)}_j(\omega)$, and $D^{(1)}(\omega) \equiv \{D^{(1)}_j(\omega)\}_j$.

A collection of intervals $\overline{I}^{(1)} \equiv \{\overline{I}^{(1)}_j\}_j$ and sets of indices $\overline{D}^{(1)} \equiv \{\overline{D}^{(1)}_j\}_j$ is called **admissible** if there exists an ω such that $I^{(1)}_j(\omega) = \overline{I}^{(1)}_j$ and $D^{(1)}_j(\omega) = \overline{D}^{(1)}_j$ for all j's. A pair $J^{(1)}_L := \{I^{(1)}_j, D^{(1)}_j\}$ is called an itinerary at level 1, or an 1-itinerary, and we denote by

$$\mathcal{I}^{(1)}_L \equiv \{J^{(1)}_L\}$$

the (finite) set of all admissible level 1-itineraries. For fixed $\overline{J}^{(1)}_L := \{\overline{I}^{(1)}_j, \overline{D}^{(1)}_j\}$ we will denote

$$\mathcal{E}(\overline{J}^{(1)}_L) = \{\omega : \ I^{(1)}(\omega) = \overline{I}^{(1)}, \ D^{(1)}(\omega) = \overline{D}^{(1)}\}.$$

1-Pedestals. Fix an admissible pair $J^{(1)}_L = \{I^{(1)}, D^{(1)}\}$ and consider all 1-blocks $I^{(1)}_j$ for which $|D^{(1)}_j| = 1$. (It implies that on $\mathcal{E}(J^{(1)}_L)$ some event $H_{i_j}(I^{(1)}_j)$ occurs.) For such 1-blocks, the set of



Figure 1: Adjustments: part *a*) shows the deterministic 1–blocks $\widetilde{I}^{(1)}_j$, bold-face segments show location of the defects. Part *b*) shows how these blocks were adjusted. $\widetilde{I}^{(1)}_5$ and $\widetilde{I}^{(1)}_6$, merge into a single 1–block $I^{(1)}_5$.

vertices $x \in I^{(1)}_j$ to the left (resp. right) of the defected 0–block in $I^{(1)}_j$ will be called *left 1-pedestal of* $I^{(1)}_j$ (resp. *right 1-pedestal*) and denoted by $\Upsilon_{\mathcal{L}}(I^{(1)}_j)$ (resp. $\Upsilon_{\mathcal{R}}(I^{(1)}_j)$). The vertices in each of these 1-pedestals are connected by open nearest neighbor edges. Define

$$\begin{aligned} G^+(I^{(1)}_j, D^{(1)}_j) &= \mathcal{E}(J^{(1)}_L) \cap [\omega : \exists x \in \Upsilon_{\mathcal{L}}(I^{(1)}_j), y \in \Upsilon_{\mathcal{R}}(I^{(1)}_j),\ y - x \le \lfloor l_1^{\alpha'/\alpha} \rfloor : \omega_{\{x,y\}} = 1], \\ B^+(I^{(1)}_j, D^{(1)}_j) &= \mathcal{E}(J^{(1)}_L) \setminus G^+(I^{(1)}_j). \end{aligned} \tag{2.3}$$

If $G^+(I^{(1)}_j, D^{(1)}_j)$ occurs, and the pair (x, y) such that $x \in \Upsilon_{\mathcal{L}}(I^{(1)}_j), y \in \Upsilon_{\mathcal{R}}(I^{(1)}_j),\ y-x \le \lfloor l_1^{\alpha'/\alpha} \rfloor$, $\omega_{\{x,y\}} = 1$ is not unique, we choose one in arbitrary way, and, once the pair (x, y) is chosen, the interval $[x + 1, y - 1]$ will be called defected part of $I^{(1)}_j$, and denoted by $\mathcal{D}(I^{(1)}_j)$. In this case we define $\Upsilon(I^{(1)}_j) = \Big(\Upsilon_{\mathcal{L}}(I^{(1)}_j) \cup \Upsilon_{\mathcal{R}}(I^{(1)}_j)\Big) \setminus \mathcal{D}(I^{(1)}_j)$. If we are in the case 3) as described above, for the leftmost interval we take $\Upsilon(I^{(1)}_j) = \Upsilon_{\mathcal{R}}(I^{(1)}_j)$ (resp. $\Upsilon(I^{(1)}_j) = \Upsilon_{\mathcal{L}}(I^{(1)}_j)$).

On the other hand, for those j such that $D^{(1)}_j = \emptyset$, we get that on $\mathcal{E}(J^{(1)}_L)$ the event $G(I^{(1)}_j)$ occurs, which means that there are no closed nearest neighbor edges in $I^{(1)}_j$, and we define the pedestal $\Upsilon(I^{(1)}_j) = I^{(1)}_j$.

This completes the first step, associating with each ω an itinerary $J^{(1)}_L(\omega) = \{I^{(1)}(\omega), D^{(1)}(\omega)\}$, and the "state" G, G^+, B or B^+ to each 1–block of the itinerary $J^{(1)}_L$. For the next step of the procedure it is irrelevant which particular sub-type G or G^+ (resp. B or B^+) occurs, without creating ambiguities, we will simply denote their occurrence by the letter G (resp. B).

In particular, a 1–pedestal $\Upsilon(I^{(1)}_j)$ is given by the vertices of open oriented path with all edges, except possibly one, being nearest neigbhour, and this larger edge has length at most $\lfloor l_1^{\alpha'/\alpha} \rfloor$. For each 1–block, except possibly the two which contain the extremes $-L$ or L, the pedestal connects left and right endpoints of the interval.

Level *k* . Let $2 \le k \le M$, and assume that we have completed the step $(k - 1)$ of the recursion. This means that for each $\omega \in \Omega_L$ and any $r = 1, \dots, k - 1$ the following objects are defined:

- the collection of r–blocks $I^{(r)}(\omega) = \{I^{(r)}_j(\omega)\}_j$, such that $\cup_j I^{(r)}_j(\omega) = [-L, L]$, and any two adjacent intervals share exactly an endpoint. Moreover, the uniform bound holds:

$$l_r - (2\lfloor l_r^{\alpha'/\alpha} \rfloor + 6l_{r-1}) < |I^{(r)}_j(\omega)| \le 3l_r + 6l_{r-1}, \tag{2.4}$$

- each of the $I_j^{(r)}(\omega)$ can be in two possible states G or B:
  If $I_j^{(r)}(\omega)$ is in the state B (or $B(I_j^{(r)}(\omega))$ occurs), we say that $I_j^{(r)}(\omega)$ is *defected*.
  If $I_j^{(r)}(\omega)$ is in the state G (or $G(I_j^{(r)}(\omega))$ occurs) and it is not the leftmost or the rightmost interval of the partition, then ω has an r–pedestal $\Upsilon(I_j^{(r)})$ given by vertices of an open oriented path from the left to the right boundary of $I_j^{(r)}(\omega)$. If $I_j^{(r)}(\omega)$ is the leftmost (resp. the rightmost) interval, an r–pedestal $\Upsilon(I_j^{(r)})$ is given by vertices of an open oriented path which starts from some vertex $x \in [-L, -L+\lfloor l_r^{\alpha'/\alpha} \rfloor]$ and ends at the right boundary of $I_j^{(r)}(\omega)$ (resp. starts from the left boundary of $I_j^{(r)}(\omega)$ and ends at some vertex $x \in [L - \lfloor l_r^{\alpha'/\alpha} \rfloor, L]$.

- the collection $D^{(r)}(\omega) = \{D_j^{(r)}(\omega)\}_j$, where $D_j^{(r)}(\omega)$ is the set of labels of the defected $(r-1)$–blocks which are contained in $I_j^{(r)}(\omega)$.

For ω fixed, the sequence of pairs

$$J_L^{(k-1)}(\omega) = \{(I^{(1)}(\omega), D^{(1)}(\omega)), \ldots, (I^{(k-1)}(\omega), D^{(k-1)}(\omega))\},$$

is called $(k-1)$–*itinerary*, and each element $(I^{(r)}, D^{(r)})$, $1 \le r \le k-1$ is called r–th step of the itinerary. For fixed $\overline{J}_L^{(k-1)} = \{(\overline{I}^{(1)}, \overline{D}^{(1)}), \ldots, (\overline{I}^{(k-1)}, \overline{D}^{(k-1)})\}$ a possible realization of a $(k-1)$–itinerary, we denote

$$\mathcal{E}(\overline{J}_L^{(k-1)}) = \{\omega : J_L^{(k-1)}(\omega) = \overline{J}_L^{(k-1)}\},$$

and we shall now see how to define the k–blocks and the continuation to a k–itinerary. When $k = M$ we will end up with only two intervals.

Construction of k - blocks. Fix $J_L^{(k-1)} = \{(I^{(1)}, D^{(1)}), \ldots, (I^{(k-1)}, D^{(k-1)})\}$ a possible realization of a $(k-1)$–itinerary. For each $z \in [-L, L]$ we set $j_z^k = \min\{j : z \in I_j^{(k-1)}\}$, $\hat{j}_i^k = j^k_{x_i^{(k)}}$, cf. notation after (2.1), $i = -l_M/l_k, \ldots, l_M/l_k - 1$, and define the intervals:

$$\widetilde{I}_i^{(k)} = \bigcup_{s=\hat{j}_i^k+1}^{\hat{j}_{i+1}^k} I_s^{(k-1)} =: [a_i^{(k)}, a_{i+1}^{(k)}]$$

as well as the following partition of $\mathcal{E}(J_L^{(k-1)})$:

$$\begin{aligned}
G(\widetilde{I}_i^{(k)}) &= \mathcal{E}(J_L^{(k-1)}) \cap \bigcap_{s=\hat{\jmath}_i^k+1}^{\hat{\jmath}_{i+1}^k} G(I_s^{(k-1)}), \\
H_s(\widetilde{I}_i^{(k)}) &= \mathcal{E}(J_L^{(k-1)}) \cap \left[B(I_s^{(k-1)}) \cap \bigcap_{u=\hat{\jmath}_i^k+1, u\neq s}^{\hat{\jmath}_{i+1}^k} G(I_u^{(k-1)}) \right], \\
H(\widetilde{I}_i^{(k)}) &= \bigcup_{s=\hat{\jmath}_i^k+1}^{\hat{\jmath}_{i+1}^k} H_s(\widetilde{I}_i^{(k)}), \\
B(\widetilde{I}_i^{(k)}) &= \mathcal{E}(J_L^{(k-1)}) \setminus \left(G(\widetilde{I}_i^{(k)}) \cup H(\widetilde{I}_i^{(k)}) \right). \qquad (2.5)
\end{aligned}$$

Adjustment. Given $\omega \in \mathcal{E}(J_L^{(k-1)})$ consider all i for which $H_s(\widetilde{I}_i^{(k)})$ occurs for s such that the distance of the defected $(k-1)$–block $I_s^{(k-1)} \subset \widetilde{I}_i^{(k)}$ to the left endpoint $a_i^{(k)}$ (right endpoint $a_{i+1}^{(k)}$, resp.) is less than $\lfloor l_k^{\alpha'/\alpha} \rfloor$. If this set is non-empty take arbitrarily any such $\widetilde{I}_i^{(k)}$.

If the selected $\widetilde{I}_i^{(k)}$ is the leftmost (resp. the rightmost) interval in $[-L, L]$, we set $I_i^{(k)} = \widetilde{I}_i^{(k)}$, and say that $G(I_i^{(k)})$ occurs (or that $I_i^{(k)}$ is in G state for this ω).

If the selected $\widetilde{I}_i^{(k)}$ is not one of these two extremal intervals, we then check if $\widetilde{I}_{i-1}^{(k)}$ (resp. $\widetilde{I}_{i+1}^{(k)}$) has a defected block $I_r^{(k-1)}$ at distance at most $3\lfloor l_{k-1}^{\alpha'/\alpha} \rfloor$ from $a_i^{(k)}$ (resp. from $a_{i+1}^{(k)}$), and

1) If yes, then we consider a new interval $I_{i-1}^{(k)} = \widetilde{I}_{i-1}^{(k)} \cup \widetilde{I}_i^{(k)}$ (respectively $I_i^{(k)} = \widetilde{I}_i^{(k)} \cup \widetilde{I}_{i+1}^{(k)}$) and say that $B(I_{i-1}^{(k)})$ (resp. $B(I_i^{(k)})$) occurs, or that the corresponding interval is in state B;

2) If not, then we consider two new intervals:

$$I_{i-1}^{(k)} = \bigcup_{s=\hat{\jmath}_{i-1}^k+1}^{j^k_{a_i^{(k)}-l_k^{\alpha'/\alpha}}-1} I_s^{(k-1)}, \qquad I_i^{(k)} = \bigcup_{s=j^k_{a_i^{(k)}-l_k^{\alpha'/\alpha}}}^{\hat{\jmath}_{i+1}^k} I_s^{(k-1)} \qquad (2.6)$$

$$\Bigg(\text{respectively,} \qquad I_i^{(k)} = \bigcup_{s=\hat{\jmath}_i^k+1}^{j^k_{a_{i+1}^{(k)}+l_k^{\alpha'/\alpha}}} I_s^{(k-1)} \qquad I_{i+1}^{(k)} = \bigcup_{s=j^k_{a_{i+1}^{(k)}+l_k^{\alpha'/\alpha}}+1}^{\hat{\jmath}_{i+2}^k} I_s^{(k-1)}\Bigg), \qquad (2.7)$$

and say that $H(I_i^{(k)})$ and $G(I_{i-1}^{(k)})$, $H(I_{i-1}^{(k)})$ or $B(I_{i-1}^{(k)})$ occurs if respectively $G(\widetilde{I}_{i-1}^{(k)})$, $H(\widetilde{I}_{i-1}^{(k)})$ or $B(\widetilde{I}_{i-1}^{(k)})$ do occur (respectively $H(I_{i-1}^{(k)})$ and $G(I_{i+1}^{(k)})$, $H(I_{i+1}^{(k)})$ or $B(I_{i+1}^{(k)})$ occurs if respectively $G(\widetilde{I}_{i+1}^{(k)})$, $H(\widetilde{I}_{i+1}^{(k)})$ or $B(\widetilde{I}_{i+1}^{(k)})$ do occur).

Finally we set $I_i^{(k)} = \widetilde{I}_i^{(k)}$ if $\widetilde{I}_i^{(k)}$ was not involved in the adjustment and say $G(I_i^{(k)})$, $H(I_i^{(k)})$ or $B(I_i^{(k)})$ occurs if respectively $G(\widetilde{I}_i^{(k)})$, $H(\widetilde{I}_i^{(k)})$ or $B(\widetilde{I}_i^{(k)})$ do occur.

If after this step we are still left with intervals $I_i^{(k)}$ for which $H(I_i^{(k)})$ occurs and its defected interval $I_s^{(k-1)}$ stays within distance $\lfloor l_k^{\alpha'/\alpha} \rfloor$ from one of the endpoints of $I_i^{(k)}$, then we repeat the above procedure. After finitely many steps of this adjustment procedure all $I_i^{(k)}$ for which $H(I_i^{(k)})$ occurs have their defected $(k-1)$–block at distance larger than $\lfloor l_k^{\alpha'/\alpha} \rfloor$ from the boundary of $I_i^{(k)}$.

Once the adjustments are completed, the final intervals are called k-blocks, and re-numerated from left to right as $I_j^{(k)},\ j = 1, \ldots$. We then consider the collection $D^{(k)} = \{D_j^{(k)}\}_j$ where $D_j^{(k)}$ gives the labels of the defected $(k-1)$–blocks contained in $I_j^{(k)}$. A sequence $(I^{(k)}, D^{(k)})$ which can be obtained in this way (for some $\omega \in \Omega_L$) is said to be **admissible** for $J_L^{(k-1)}$, in which case $J_L^{(k)} := \{(I^{(s)}, D^{(s)}),\ 1 \le s \le k\}$ is a k–itinerary.

By $\mathcal{I}^{(k)}(J_L^{(s)})$ we denote the collection of all k–itineraries that are continuation of $J_L^{(k-1)}$.

We can always write $I_j^{(k)} = \bigcup_{s_0(j)}^{s_1(j)} I_s^{(k-1)}$, and simple recursive estimate holds:

$$l_k - (2\lfloor l_k^{\alpha'/\alpha} \rfloor + 6 l_{k-1}) < |I_j^{(k)}| \le 3 l_k + 6 l_{k-1}. \tag{2.8}$$

k-Pedestals. For ω fixed, let $I_j^{(k)}(\omega)$ be a k–block for which after adjustment $G(I_j^{(k)})$ or $H(I_j^{(k)})$ occurs. For such k–blocks we will define k–pedestals.

- if $G(I_j^{(k)})$ occurs and if $I_j^{(k)}(\omega)$ is not leftmost (resp. rightmost) interval in $[-L, L]$, then all its sub-blocks $I_s^{(k-1)}$ are in G state, and we define $\Upsilon(I_j^{(k)}) = \bigcup_{s_0(j)}^{s_1(j)} \Upsilon(I_s^{(k-1)})$.

- if $G(I_j^{(k)})$ occurs and $I_j^{(k)}(\omega)$ is the leftmost (resp. rightmost) interval in $[-L, L]$ which contains unique defected $(k-1)$–block $I_r^{(k-1)}$ within distance $\lfloor l_k^{\alpha'/\alpha} \rfloor$ from its left (resp. right) boundary then $\Upsilon(I_j^{(k)}) = \bigcup_{r+1}^{s_1(j)} \Upsilon(I_s^{(k-1)})$ (resp. $\Upsilon(I_j^{(k)}) = \bigcup_{s_0(j)}^{r-1} \Upsilon(I_s^{(k-1)})$).

- if $H_r(I_j^{(k)})$ occurs[4], we define $\Upsilon_{\mathcal{L}}(I_j^{(k)}) = \bigcup_{s_0(j)}^{r-1} \Upsilon(I_s^{(k-1)})$ and $\Upsilon_{\mathcal{R}}(I_j^{(k)}) = \bigcup_{r+1}^{s_1(j)} \Upsilon(I_s^{(k-1)})$, called left and right pedestals[5] of $I_j^{(k)}$, and check if there exists $x \in \Upsilon_{\mathcal{L}}(I_j^{(k)})$ and $y \in \Upsilon_{\mathcal{R}}(I_j^{(k)})$ with $y - x \le \lfloor l_k^{\alpha'/\alpha} \rfloor$ such that $\omega_{\{x,y\}} = 1$:

  – If yes, we say that $I_j^{(k)}$ is in G state, and if the pair (x, y) with $x \in \Upsilon_{\mathcal{L}}(I_j^{(k)}), y \in \Upsilon_{\mathcal{R}}(I_j^{(k)}),\ y - x \le \lfloor l_k^{\alpha'/\alpha} \rfloor$, and $\omega_{\{x,y\}} = 1$ is not unique, we choose one in an arbitrary way, and, once (x, y) is chosen, denote $\mathcal{D}(I_j^{(k)}) = [x+1, y-1]$, and define

  $$\Upsilon(I_j^{(k)}) = \Big(\Upsilon_{\mathcal{L}}(I_j^{(k)}) \cup \Upsilon_{\mathcal{R}}(I_j^{(k)})\Big) \setminus \mathcal{D}(I_j^{(k)}).$$

  – If such an open edge $\{x, y\}$ does not exist we say that $I_j^{(k)}$ is in B state.

---

[4] i.e. $I_r^{(k-1)}$ is the only defected block within $I_j^{(k)}$

[5] From the occurrence of $G(I_s^{(k-1)})$ for all other $(k-1)$–blocks within $I_j^{(k)}$, we know there exists an open oriented path connecting the left boundary of $I_j^{(k)}$ to the right boundary of $I_{r-1}^{(k-1)}$ and an open oriented path connecting the left boundary of $I_{r+1}^{(k-1)}$ to the right boundary of $I_j^{(k)}$. These paths are obtained by concatenation of the corresponding $\Upsilon(I_s^{(k-1)}),\ s_0(j) \le s \le r-1$ and $r+1 \le s \le s_1(j)$, respectively.


a)
$I_j^{(2)}$
$I_1^{(1)}$ $I_2^{(1)}$ $I_3^{(1)}$ $I_4^{(1)}$ $I_5^{(1)}$ $I_6^{(1)}$ $I_7^{(1)}$ $I_8^{(1)}$ $I_9^{(1)}$ $I_{10}^{(1)}$
b)
$I_j^{(2)}$
x_1 y_1 x_2 y_2 x_4 y_4 x_7 y_7 x_8 y_8 x_{10} y_{10}
$I_1^{(1)}$ $I_2^{(1)}$ $I_3^{(1)}$ $I_4^{(1)}$ $I_5^{(1)}$ $I_6^{(1)}$ $I_7^{(1)}$ $I_8^{(1)}$ $I_9^{(1)}$ $I_{10}^{(1)}$
c)
$I_j^{(2)}$
x_j y_j

Figure 2: Pedestals and defects: part a) shows the deterministic 1–blocks $I_i^{(1)}$, $1 \leq i \leq 10$, located in the 2–block $I_j^{(2)}$; bold-face segments show location of the 0–level defects. Part b) shows construction of 1–level pedestals, marked by light-gray strips. The block $I_6^{(1)}$ is a defected 1–block. The segments (x_i, y_i) are "enlarged" defects in $I_i^{(1)}$. Part c) shows creation of 2–level pedestals, marked by dark-gray strips, concatenated by long range edges. The segment (x_j, y_j) is enlarged defect for $I_j^{(2)}$.

In other words, we consider the event

$$G^+(I_j^{(k)}) \;\; = \;\; H(I_j^{(k)}) \cap [\omega : \exists x \in \Upsilon_{\mathcal{L}}(I_j^{(k)}), y \in \Upsilon_{\mathcal{R}}(I_j^{(k)}),\ y - x \leq \lfloor l_k^{\alpha'/\alpha} \rfloor : \omega_{\{x,y\}} = 1],$$

whose occurrence implies the k–block to be in G state, i.e. when dealing at larger scales we do not distinguish if $I_j^{(k)}$ is in G state due to the occurrence of $G(I_j^{(k)})$ or $G^+(I_j^{(k)})$.

This completes the k–th step, associating with each itinerary $J^{(k-1)}$ its continuation with a sequence of k–blocks $I^{(k)} = \{I_j^{(k)}\}_j$, re-numerated from left to right. Moreover, with each k–block we associate one of the states G or B.

Structure of pedestals. First we state a simple geometric property of pedestals, which will be used in estimating the conditional probability of the event G^+ defined above, given H. Our goal is to show that if a k–block, $k \geq 1$, $I^{(k)} = [s, s']$ contains only one defected $(k-1)$–block, here denoted by $[a, a']$, and there are corresponding left and right pedestals $\Upsilon_{\mathcal{L}}$ and $\Upsilon_{\mathcal{R}}$, spanning from s to a and from a' to s', respectively, then there exists a constant $C \equiv C(\alpha, \alpha')$ such that

$$\left| \Upsilon_{\mathcal{L}} \cap [a - \lfloor l_k^{\alpha'/\alpha} \rfloor,\, a] \right| \geq C l_k^{\alpha'/\alpha} \quad \text{and} \quad \left| \Upsilon_{\mathcal{R}} \cap [a',\, a' + \lfloor l_k^{\alpha'/\alpha} \rfloor] \right| \geq C l_k^{\alpha'/\alpha}. \tag{2.9}$$

Inequality (2.9) follows trivially from the following recursive relation: if we have a k–block $I^{(k)} = \cup_{s_0}^{s_1} I_s^{(k-1)}$ which is in G state, then

$$\left| \Upsilon(I^{(k)}) \right| \geq \sum_{s:\, [G(I_s^{(k-1)}) \text{ occurs}]} \left| \Upsilon(I_s^{(k-1)}) \right| - l_k^{\alpha'/\alpha}.$$

We will end this section with the estimate which will be important for the recursive step in the next section, and after that we also fix the parameters which will determine the choice of p close to one, as in (1.5). Recall that by $H_z(I_j^{(k)}) \subset H(I_j^{(k)})$ we denote the event, for which the unique defected $(k-1)$–block within $I_j^{(k)}$ has index z, and by construction stays at distance larger than $\lfloor l_k^{\alpha'/\alpha} \rfloor$ from the boundaries of $I_j^{(k)}$.

Lemma 2.1 *There exists* $\eta \equiv \eta(\alpha, \alpha', l_1)$ *with* $\eta \searrow 0$ *as* $l_1 \nearrow +\infty$ *and such that the following estimate for the conditional probability with respect to the product measure (defined right above (1.1))*

$$\nu\big[\exists x \in \Upsilon_{\mathcal{L}}(I_j^{(k)}), y \in \Upsilon_{\mathcal{R}}(I_j^{(k)}),\ y - x \le \lfloor l_k^{\alpha'/\alpha} \rfloor : \omega_{\{x,y\}} = 1 \big|\, H(I_j^{(k)})\big] \ge 1 - l_{k-1}^{-\beta(1-\eta)(\alpha'-1)} \tag{2.10}$$

holds for $k \ge 2$. *For* $k = 1$ *the r.h.s in (2.10) is replaced by* $1 - l_1^{-\beta(1-\eta)(\alpha'-1)/\alpha}$.

Proof. We show the above estimate by conditioning on $H_z(I_j^{(k)})$, uniformly in z, and we make repeated use of the following upper and lower bounds: if I and I' are two intervals, and $3 \le d = \mathrm{dist}(I, I')$, then

$$C^-\ J(I, I') \le \sum_{\substack{x \in I \cap \mathbb{Z} \\ y \in I' \cap \mathbb{Z}}} \frac{1}{|x-y|^2} \le C^+\ J(I, I'), \tag{2.11}$$

holds with $C^\pm = (1 \pm 2/d)^2$ and

$$J(I, I') = \int_{I \times I'} dx\, dy \frac{1}{|x-y|^2} = \ln \frac{(|I| + d)(|I'| + d)}{d\,(|I| + |I'| + d)}. \tag{2.12}$$

Notice that we have $C^-\,(|x-y|-2)^{-2} \le |x-y|^{-2} \le C^+\,(|x-y|+2)^{-2}$ for $|x-y| \ge d$. We shall need also the inequality

$$J(I, I') \le 4 \frac{|I'|}{|I''|} J\,(I, I'') \tag{2.13}$$

which holds for every I, I' and I'' such that $I' \subset I''$ and $d' = \mathrm{dist}(I, I'') \ge |I''|$. Indeed, setting $f(x) = \int_I dy\, |x-y|^{-2}$, for $x \in I''$, straightforward calculations give that under the above conditions:

$$f(x') \le 4 f(x'') \quad \text{for each} \quad x' \in I', x'' \in I''$$

from where the inequality (2.13) follows upon integration.

If $k \ge 2$ and $H_z(I_j^{(k)})$ occurs, we have the left k–pedestal $\Upsilon_{\mathcal{L}}(I_j^{(k)})$ spanning from the left endpoint of $I_j^{(k)}$ to the left endpoint of $I_z^{(k-1)}$, and the right k-pedestal $\Upsilon_{\mathcal{R}}(I_j^{(k)})$, spanning from the right endpoint of $I_z^{(k-1)}$ to the right endpoint of $I_j^{(k)}$. Take two segments $S_z^{\mathcal{L}}$ and $S_z^{\mathcal{R}}$, such that $|S_z^{\mathcal{L}}| = |S_z^{\mathcal{R}}| = \lfloor l_k^{\alpha'/\alpha} \rfloor / 3$, lying immediately to the left and, respectively, to the right of $I_z^{(k-1)}$. Denote

$$\begin{aligned} \widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}) &= \Upsilon_{\mathcal{L}}(I_j^{(k)}) \cap S_z^{\mathcal{L}}, \\ \widehat{\Upsilon}_{\mathcal{R}}(I_j^{(k)}) &= \Upsilon_{\mathcal{R}}(I_j^{(k)}) \cap S_z^{\mathcal{R}}. \end{aligned}$$

Then

$$\nu[\text{all edges } \{x,y\},\ x\in\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}),\ y\in\widehat{\Upsilon}_{\mathcal{R}}(I_j^{(k)}) \text{ are closed}]$$
$$\leq \prod_{\substack{x\in S_z^{\mathcal{L}}\\ y\in S_z^{\mathcal{R}}}} q_{\{x,y\}} \prod_{\substack{x\in S_z^{\mathcal{L}}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)})\\ y\in S_z^{\mathcal{R}}}} q_{\{x,y\}}^{-1} \prod_{\substack{x\in S_z^{\mathcal{L}}\\ y\in S_z^{\mathcal{R}}\setminus\widehat{\Upsilon}_{\mathcal{R}}(I_j^{(k)})}} q_{\{x,y\}}^{-1}.$$

Applying (2.11) to $S_z^{\mathcal{L}}$ and $S_z^{\mathcal{R}}$ we immediately get the following bound:

$$\prod_{\substack{x\in S_z^{\mathcal{L}}\\ y\in S_z^{\mathcal{R}}}} q_{\{x,y\}} = \exp\Big\{-\sum_{\substack{x\in S_z^{\mathcal{L}}\\ y\in S_z^{\mathcal{R}}}} \frac{\beta}{|x-y|^2}\Big\} \leq l_{k-1}^{-\beta(\alpha'-1)(1-b)}, \tag{2.14}$$

where $b\equiv b(\alpha',l_1)$ and $b\searrow 0$ when $l_1\nearrow+\infty$. The same computation gives that if a 1-block I has a unique closed edge $\{a,a+1\}$ with both $a,a+1$ at distance larger than $l_1^{\alpha'/\alpha}$ from the endpoints of I, then the probability that there is an open edge $\{x,y\}$ with $x<a<y$, $y-x\leq l_1^{\alpha'/\alpha}$ is larger than or equal of $1-l_1^{-\beta(1-\eta)(\alpha'-1)/\alpha}$.

On the other hand denoting by $\mathcal{D}_n(S_z^{\mathcal{L}}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}))$, $0\leq n\leq k-2$ (resp. $\mathcal{D}_n(S_z^{\mathcal{R}}\setminus\widehat{\Upsilon}_{\mathcal{R}}(I_j^{(k)}))$) the set of vertices which belong to all n–level defects contained in the segment $S_z^{\mathcal{L}}$ (resp. $S_z^{\mathcal{R}}$), we get

$$\prod_{\substack{x\in S_z^{\mathcal{L}}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)})\\ y\in S_z^{\mathcal{R}}}} q_{\{x,y\}} = \prod_{n=0}^{k-2}\prod_{\substack{x\in\mathcal{D}_n(S_z^{\mathcal{L}}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}))\\ y\in S_z^{\mathcal{R}}}} q_{\{x,y\}} = \exp\Big\{-\sum_{n=0}^{k-2}\sum_{\substack{x\in\mathcal{D}_n(S_z^{\mathcal{L}}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}))\\ y\in S_z^{\mathcal{R}}}} \frac{\beta}{|x-y|^2}\Big\}.$$

Once again, applying (2.11) for each $0\leq n\leq k-2$ and taking into account the structure of n–level pedestals together with (2.13), we have

$$\begin{aligned}\sum_{n=0}^{k-2}\sum_{\substack{x\in\mathcal{D}_n(S_z^{\mathcal{L}}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}))\\ y\in S_z^{\mathcal{R}}}} \frac{\beta}{|x-y|^2} &\leq C^+\sum_{n=0}^{k-2}\sum_{\nu} J\left(I'_\nu, I^R\right)\\ &\leq C\sum_{n=0}^{k-2}\frac{l_{n+1}^{\alpha'/\alpha}}{l_{n+1}}\sum_{\nu} J\left(I''_\nu, I^R\right)\\ &\leq C' l_1^{\alpha'/\alpha-1} J\left(I^{\mathcal{L}}, I^{\mathcal{R}}\right)\end{aligned}$$

where $I^{\mathcal{L}(\mathcal{R})}$, I'_ν and I''_ν are intervals in $\mathbb{R}$ such that $I^{\mathcal{L}(\mathcal{R})}\cap\mathbb{Z}=S_z^{\mathcal{L}(\mathcal{R})}$, $\bigcup_\nu(I'_\nu\cap\mathbb{Z})=\mathcal{D}_n(S_z^{L}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}))$ and the sum $\sum_\nu$ is taken over all indices ν of $(n+1)$–blocks $I_\nu^{(n+1)}=I''_\nu\cap\mathbb{Z}$, where the defected n–blocks are located. The condition to apply (2.13) in the first inequality above follows from $3l_{k-1}+6l_{k-2}\leq l_k^{\alpha'/\alpha}$ which is true for any $k\geq 2$, provided l_1 has geen taken large enough. From this we can easily get that

$$\prod_{\substack{x\in S_z^{\mathcal{L}}\setminus\widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)})\\ y\in S_z^{\mathcal{R}}}} q_{\{x,y\}} \geq l_{k-1}^{-\beta(\alpha'-1)b'}, \tag{2.15}$$

where $b' \equiv b'(\alpha, \alpha', l_1)$ and $b' \searrow 0$ when $l_1 \nearrow +\infty$. Analogous bound holds for the third term. Finally, from the upper bound for the length of a $(k-1)$-block, we have

$$\begin{aligned} & [\omega : \exists x \in \Upsilon_{\mathcal{L}}(I_j^{(k)}), y \in \Upsilon_{\mathcal{R}}(I_j^{(k)}),\ y - x \le \lfloor l_k^{\alpha'/\alpha} \rfloor : \omega_{\{x,y\}} = 1]^c \\ \subseteq\ & [\text{all edges } \{x,y\},\ x \in \widehat{\Upsilon}_{\mathcal{L}}(I_j^{(k)}),\ y \in \widehat{\Upsilon}_{\mathcal{R}}(I_j^{(k)}) \text{ are closed}], \end{aligned} \tag{2.16}$$

the statement of the Lemma follows from (2.14) and (2.15). $\square$

Fixing the parameters. For fixed $\beta > 1$, which is the first main parameter of the model we choose the pair $\alpha,\ \alpha'$ with $1 < \alpha' < \alpha < 2$ such that

$$\beta(\alpha' - 1) - \frac{2(\alpha - 1)^2}{2 - \alpha} > \alpha - 1, \tag{2.17}$$

i.e. $\beta(\alpha' - 1) > \alpha\,(\alpha - 1)\,/(2 - \alpha)$. Now we fix $l_1 > 1$ so large that the parameter $\eta = \eta(\alpha, \alpha', l_1)$ in (2.10) becomes so close to zero, that

$$\beta(1 - \eta)(\alpha' - 1) - \frac{2(\alpha - 1)^2}{2 - \alpha} > \alpha - 1. \tag{2.18}$$

Finally we fix

$$\delta > \frac{2(\alpha - 1)}{2 - \alpha} \tag{2.19}$$

such that

$$\beta(1 - \eta)(\alpha' - 1) - \delta(\alpha - 1) > \alpha - 1. \tag{2.20}$$

Inequalities (2.10), (2.19) and (2.20) will be crucial for the inductive estimates.

3 Decomposition of the product measure. Basic estimates

Let ν_L be the product measure on Ω_L, the space of configurations over edges with both end-vertices in $[-L, L]$. Looking at successive projections of this measure, we need a notation that emphasize the set of edges which have been integrated out. This explains the somehow heavy notation below.

Proposition 3.1 *For $L = l_M$ and $k = 0, \ldots, M$, the measure ν_L can be written as*

$$\nu_L = \sum_{J_L^{(k)} \in \mathcal{I}_L^{(k)}} c(J_L^{(k)}) \prod_{I_i^{(k)} \in\ I^{(k)}} \Big(\gamma_{I_i^{(k)}} + \Delta_{I_i^{(k)}}\Big)\nu^o_{J_L^{(k)}}, \tag{3.1}$$

where

- *the sum $\sum_{J_L^{(k)} \in \mathcal{I}^{(k)}}$ is taken over all possible itineraries $J_L^{(k)} = \{\{I^{(i)}, D^{(i)}\}_{i=0}^{k}\}$ of length k;*

the product $\prod_{I_i^{(k)} \in I^{(k)}}$ is taken over all k–blocks $I_i^{(k)}$ which form partition of L at the last step $I^{(k)}$of the itinerary $J_L^{(k)}$.

- *The measure* $\nu^o_{J_L^{(k)}}$ *is the marginal of* ν_L*, after integrating out all variables corresponding to the edges* $e \in \cup_i \, \mathbb{E}(I_i^{(k)})$*,* $I_i^{(k)} \in I^{(k)}$*.*
- $c(J_L^{(k)}) \geq 0, \quad \sum_{J_L^{(k)} \in \mathcal{I}_L^{(k)}} c(J_L^{(k)}) = 1;$
- $\gamma_{I_i^{(k)}}$ *and* $\Delta_{I_i^{(k)}}$ *are non-negative functions satisfying the following properties:*

1)

$$\gamma_{I_i^{(k)}}(\omega)\Delta_{I_i^{(k)}}(\omega) = 0. \tag{3.2}$$

2) If $\Delta_{I_i^{(k)}}(\omega) > 0$*, in which case we say that* $I_i^{(k)}$ *is defected for* ω*, then there is no* k*–pedestal connecting the end-vertices of* $I_i^{(k)}$*. Moreover,*

$$\sup_\omega \kappa^{\mathcal{C}_{I_i^{(k)}}(\omega)} \Delta_{I_i^{(k)}}(\omega) \leq l_k^{-\delta}. \tag{3.3}$$

3) If $\gamma_{I_i^{(k)}}(\omega) > 0$*, the interval* $I_i^{(k)}$ *contains a* k*–pedestal which, unless* $I_i^{(k)}$ *is the rightmost or the leftmost for* $I^{(k)}$*, connects its end-vertices; otherwise, the* k*–pedestal is as explained in Sect. 2. Moreover, for all configurations* ω *with at least one closed nearest neighbour edge in* $I_i^{(k)}$*, one has*

$$\kappa^{\mathcal{C}_{I_i^{(k)}}(\omega)} \gamma_{I_i^{(k)}}(\omega) \leq 1. \tag{3.4}$$

When all nearest neighbour edges in $I_i^{(k)}$ *are open we only have the trivial bound* $\gamma_{I_i^{(k)}}(\omega) \leq 1$*.*

Proof. Through this section we will repeatedly use the following straightforward identity: Let $N > 0$, $c_N = 2^{1/N} - 1$, and $\{\gamma_i\}_1^N$ and $\{\Delta_i\}_1^N$ any two sets of random variables. Then

$$\prod_{j=1}^{N} (\gamma_i + \Delta_i) = \sum_{\substack{J \subset \{1,\dots,N\}: \\ J \neq \emptyset}} c_N^{|J|} \left(\prod_{j=1}^{N} \gamma_i + c_N^{-|J|} \prod_{k \in J^c} \gamma_k \prod_{j \in J} \Delta_j \right), \tag{3.5}$$

$$\sum_{\substack{J \subset \{1,\dots,N\}: \\ J \neq \emptyset}} c_N^{|J|} = 1, \qquad\qquad c_N^{-1} \leq N / \ln 2. \tag{3.6}$$

Step 1. $\boldsymbol{k = 0}$. In this case $\mathcal{I}_L^{(0)}$ has only one element $J_L^{(0)} = \{I^{(0)}, D^{(0)}\}$, and $I^{(0)} = \{I_i^{(0)}\}_{i=-L}^{L-1}$, $D^{(0)} = \{D_i^{(0)}\}_{i=-L}^{L-1}$ with $I_i^{(0)} = [i, i+1]$, and $D_i^{(0)} = \emptyset$ for all i. Thus we trivially have:

$$\nu_L = \prod_{I_i^{(0)} \in I^{(0)}} \left(p \, \mathbb{I}_{G(I_i^{(0)})} + (1-p) \mathbb{I}_{B(I_i^{(0)})} \right) \nu^o_{J_L^{(0)}} = \prod_{I_i^{(0)} \in I^{(0)}} \left(\gamma_{I_i^{(0)}} + \Delta_{I_i^{(0)}} \right) \nu^o_{J_L^{(0)}}, \tag{3.7}$$

where $\gamma_{I_i^{(0)}} = p \, \mathbb{I}_{G(I_i^{(0)})}$ and $\Delta_{I_i^{(0)}} = (1-p) \mathbb{I}_{B(I_i^{(0)})}$. For ω such that $\mathbb{I}_{B(I_i^{(0)})}(\omega) = 1$ we have $\mathcal{C}_{I_i^{(0)}}(\omega) = 2$, since in this case in $I_i^{(0)}$ the nearest neighbor edge is closed and we have two end-vertices as two disjoint connected components. Thus, by our choice of p in (1.5), we have

$$\sup_\omega \kappa^2 \Delta_{I_i^{(0)}}(\omega) \leq \frac{(\ln 2)^5}{128} l_1^{-\delta - 1} \ . \tag{3.8}$$

Item 3) is trivially checked when $k = 0$.

Proof of the recursive step. Though simpler, the procedure involved to get the estimates for $k = 1$ has minor differences with respect to the case $k \geq 2$. We point them out in the proof.

Let $k \geq 1$, and assume that we have the decomposition at step $(k-1)$, satisfying the above described properties:

$$\nu_L = \sum_{J_L^{(k-1)} \in \mathcal{I}_L^{(k-1)}} c(J_L^{(k-1)}) \prod_{I_u^{(k-1)} \in \; I^{(k-1)}} \Big(\gamma_{_{I_u^{(k-1)}}} + \Delta_{_{I_u^{(k-1)}}} \Big) \nu^o_{J_L^{(k-1)}}. \tag{3.9}$$

To obtain the decomposition and the estimates for the k - th step we proceed as follows: first, for each fixed itinerary $J^{(k-1)} \in \mathcal{I}^{(k-1)}$ we rewrite

$$\prod_{I_u^{(k-1)} \in \; I^{(k-1)}} \Big(\gamma_{_{I_u^{(k-1)}}} + \Delta_{_{I_u^{(k-1)}}} \Big) \tag{3.10}$$

as a convex sum of product-type expressions, where the sum will be taken over all possible choices of $I^{(k)}$ admissible from $I^{(k-1)}$ as the k–th step of the itinerary which starts from the last step of $J_L^{(k-1)}$. The products will be taken over all such k–blocks $I_i^{(k)} \in I^{(k)}$.

Recalling the construction of k–blocks in Sect. 2, and the notation $\hat{\jmath}_i^k = \min\{s \colon x_i^{(k)} \in I_s^{(k-1)}\}$, we write

$$\prod_{I_u^{(k-1)} \in I^{(k-1)}} \Big(\gamma_{_{I_u^{(k-1)}}} + \Delta_{_{I_u^{(k-1)}}} \Big) = \prod_{i\colon \, x_i^{(k)} \in [-L,L]} \; \prod_{s=\hat{\jmath}_i^k+1}^{\hat{\jmath}_{i+1}^k} \Big(\gamma_{_{I_s^{(k-1)}}} + \Delta_{_{I_s^{(k-1)}}} \Big). \tag{3.11}$$

For each i in the product, take $D_i^{(k)} \subseteq \{\hat{\jmath}_i^k + 1, \ldots, \hat{\jmath}_{i+1}^k\}$, $D_i^{(k)} \neq \emptyset$, and using the identity (3.5) we rewrite r.h.s. of (3.11) as

$$\sum_{\{D_i^{(k)}\}_i} c(\{D_i^{(k)}\}_i) \prod_i \Big(\prod_{s=\hat{\jmath}_i^k+1}^{\hat{\jmath}_{i+1}^k} \gamma_{_{I_s^{(k-1)}}} + c^{-|D_i^{(k)}|}_{\hat{\jmath}_{i+1}^k - \hat{\jmath}_i^k - 1} \prod_{s' \in [\hat{\jmath}_i^k+1, \hat{\jmath}_{i+1}^k] \setminus D_i^{(k)}} \gamma_{_{I_{s'}^{(k-1)}}} \prod_{s'' \in D_i^{(k)}} \Delta_{_{I_{s''}^{(k-1)}}} \Big), \tag{3.12}$$

where the sum $\sum_{\{D_i^{(k)}\}_i}$ is taken over all possible collections $\{D_i^{(k)}\}_i$, $i \in \{-l_M/l_k, \ldots, l_M/l_k - 1\}$, and the product $\prod_i$ is taken over all (not-yet-modified) intervals $\tilde{I}_i^{(k)} = \cup_{s=\hat{\jmath}_i^k+1}^{\hat{\jmath}_{i+1}^k} I_s^{(k-1)} \equiv [a_i^{(k)}, a_{i+1}^{(k)}]$. Moreover

$$c(\{D_i^{(k)}\}_i) \geq 0, \qquad \sum_{\{D_i^{(k)}\}_i} c(\{D_i^{(k)}\}_i) = 1,$$

and $D_i^{(k)}$ is the set of indices of the defected $(k-1)$–blocks $I_s^{(k-1)}$ in $\tilde{I}_i^{(k)}$.

Rearrangement of factors at step k. Recall that at this stage we are working with the fixed itinerary $J_L^{(k-1)}$, and now we will analyze each term in (3.12). Thus $\{D_i^{(k)}\}_i$ is fixed, and we rearrange each product

$$\prod_i \Big(\prod_{s=\hat{\jmath}_i^k+1}^{\hat{\jmath}_{i+1}^k} \gamma_{_{I_s^{(k-1)}}} + c^{-|D_i^{(k)}|}_{\hat{\jmath}_{i+1}^k - \hat{\jmath}_i^k - 1} \prod_{s' \in [\hat{\jmath}_i^k+1, \hat{\jmath}_{i+1}^k] \setminus D_i^{(k)}} \gamma_{_{I_{s'}^{(k-1)}}} \prod_{s'' \in D_i^{(k)}} \Delta_{_{I_{s''}^{(k-1)}}} \Big), \tag{3.13}$$

according to locations of the defected $(k-1)$–blocks $I_s^{(k-1)},\ s \in D_i^{(k)}$ in $\tilde{I}_i^{(k)} = [a_i^{(k)}, a_{i+1}^{(k)}]$.

i) Boundary shift case. We check, as explained in Section 2, if there is an i so that $|D_i^{(k)}| = 1$, say $D_i^{(k)} = \{s\}$, and the corresponding defected $I_s^{(k-1)}$ stays at distance not more than $l_k^{\alpha'/\alpha}$ from an end-vertex of $\tilde{I}_i^{(k)}$, say $a_i^{(k)}$. For each such i, we check if $D_{i-1}^{(k)}$ contains an index v so that the corresponding $I_v^{(k-1)}$ stays at distance not more than $3l_k^{\alpha'/\alpha}$ from $a_i^{(k)}$. If it does not, then we shift the boundary between the blocks $\tilde{I}_{i-1}^{(k)}$ and $\tilde{I}_i^{(k)}$, as in (2.6). In agreement, we rewrite the product of the two corresponding factors in (3.12)

$$\Big(\prod_{s=\hat{\jmath}_{i-1}^k+1}^{\hat{\jmath}_i^k} \gamma_{I_s^{(k-1)}} + c_{\hat{\jmath}_i^k - \hat{\jmath}_{i-1}^k - 1}^{-|D_{i-1}^{(k)}|} \prod_{s' \in [\hat{\jmath}_{i-1}^k+1, \hat{\jmath}_i^k] \setminus D_{i-1}^{(k)}} \gamma_{I_{s'}^{(k-1)}} \prod_{s'' \in D_{i-1}^{(k)}} \Delta_{I_{s''}^{(k-1)}} \Big)$$

$$\times \ \Big(\prod_{s=\hat{\jmath}_i^k+1}^{\hat{\jmath}_{i+1}^k} \gamma_{I_s^{(k-1)}} + c_{\hat{\jmath}_{i+1}^k - \hat{\jmath}_i^k - 1}^{-|D_i^{(k)}|} \prod_{s' \in [\hat{\jmath}_i^k+1, \hat{\jmath}_{i+1}^k] \setminus D_i^{(k)}} \gamma_{I_{s'}^{(k-1)}} \prod_{s'' \in D_i^{(k)}} \Delta_{I_{s''}^{(k-1)}} \Big) \tag{3.14}$$

as (our condition implies, in particular that $D_{i-1}^{(k)} \cap [j^k_{a_i^{(k)} - \lfloor l_k^{\alpha'/\alpha} \rfloor} + 1, \hat{\jmath}_i^k] = \emptyset$):

$$\Big(\prod_{s=\hat{\jmath}_{i-1}^k+1}^{j^k_{a_i^{(k)} - \lfloor l_k^{\alpha'/\alpha} \rfloor}} \gamma_{I_s^{(k-1)}} + c_{j_i^k - j_{i-1}^k - 1}^{-|D_{i-1}^{(k)}|} \prod_{s' \in [j_{i-1}^k+1, j^k_{a_i^{(k)} - \lfloor l_k^{\alpha'/\alpha} \rfloor}] \setminus D_{i-1}^{(k)}} \gamma_{I_{s'}^{(k-1)}} \prod_{s'' \in D_{i-1}^{(k)}} \Delta_{I_{s''}^{(k-1)}} \Big)$$

$$\times \ \Big(\prod_{s=j^k_{a_i^{(k)} - \lfloor l_k^{\alpha'/\alpha} \rfloor}+1}^{\hat{\jmath}_{i+1}^k} \gamma_{I_s^{(k-1)}} + c_{j_{i+1}^k - j_i^k - 1}^{-|D_i^{(k)}|} \prod_{s' \in [j^k_{a_i^{(k)} - \lfloor l_k^{\alpha'/\alpha} \rfloor}+1, \hat{\jmath}_{i+1}^k] \setminus D_i^{(k)}} \gamma_{I_{s'}^{(k-1)}} \prod_{s'' \in D_i^{(k)}} \Delta_{I_{s''}^{(k-1)}} \Big). \tag{3.15}$$

Analogous rewriting is performed in the case the defect stays close to the right boundary of $\tilde{I}_i^{(k)}$ and $\tilde{I}_{i+1}^{(k)}$ has no defect within distance $3l_k^{\alpha'/\alpha}$ from its left boundary.

ii) Two blocks merge case. Consider only the indices i for which $|D_i^{(k)}| = 1$ and the corresponding defected $(k-1)$–block stays at distance not more than $l_k^{\alpha'/\alpha}$ from an end-vertex of $\tilde{I}_i^{(k)}$, say $a_i^{(k)}$. Assume now that differently from the previous case i) the block $\tilde{I}_{i-1}^{(k)}$ has a defected $(k-1)$–block within distance $3l_k^{\alpha'/\alpha}$ from $a_i^{(k)}$. As explained in Section 2, in this situation we can not shift the boundary between the two blocks, and instead we will consider them together, as one defected block at level k. In this case the product of two terms, as in (3.14), but now corresponding to two merging blocks, is rewritten using (3.5) for a two factor product:

$$
\begin{aligned}
& c_2\Big(\prod_{s=\hat{\jmath}^k_{i-1}+1}^{\hat{\jmath}^k_{i+1}} \gamma_{I_s^{(k-1)}} + c_2^{-1} c_{\hat{\jmath}^k_i-\hat{\jmath}^k_{i-1}-1}^{-|D^{(k)}_{i-1}|} \prod_{s'\in[\hat{\jmath}^k_i+1,\hat{\jmath}^k_{i+1}]\setminus D^{(k)}_i} \gamma_{I_{s'}^{(k-1)}} \prod_{s''\in D^{(k)}_i} \Delta_{I_{s''}^{(k-1)}} \prod_{s=\hat{\jmath}^k_{i-1}+1}^{\hat{\jmath}^k_{i+1}} \gamma_{I_s^{(k-1)}} \Big) \\
+ \;\; & c_2\Big(\prod_{s=\hat{\jmath}^k_{i-1}+1}^{\hat{\jmath}^k_{i+1}} \gamma_{I_s^{(k-1)}} + c_2^{-1} c_{\hat{\jmath}^k_{i+1}-\hat{\jmath}^k_{i}-1}^{-|D^{(k)}_{i}|} \prod_{s=\hat{\jmath}^k_{i-1}+1}^{\hat{\jmath}^k_{i}} \gamma_{I_s^{(k-1)}} \prod_{s'\in[\hat{\jmath}^k_{i-1}+1,\hat{\jmath}^k_{i}]\setminus D^{(k)}_{i-1}} \gamma_{I_{s'}^{(k-1)}} \prod_{s''\in D^{(k)}_{i-1}} \Delta_{I_{s''}^{(k-1)}} \Big) \\
+ \;\; & c_2^2\Big(\prod_{s=\hat{\jmath}^k_{i-1}+1}^{\hat{\jmath}^k_{i+1}} \gamma_{I_s^{(k-1)}} + c_2^{-2} c_{\hat{\jmath}^k_{i+1}-\hat{\jmath}^k_{i}-1}^{-|D^{(k)}_{i}|} c_{\hat{\jmath}^k_i-\hat{\jmath}^k_{i-1}-1}^{-|D^{(k)}_{i-1}|} \prod_{s'\in[\hat{\jmath}^k_i+1,\hat{\jmath}^k_{i+1}]\setminus D^{(k)}_i} \gamma_{I_{s'}^{(k-1)}} \prod_{s''\in D^{(k)}_i} \Delta_{I_{s''}^{(k-1)}} \\
\times \;\; & \prod_{s'\in[\hat{\jmath}^k_{i-1}+1,\hat{\jmath}^k_{i}]\setminus D^{(k)}_{i-1}} \gamma_{I_{s'}^{(k-1)}} \prod_{s''\in D^{(k)}_{i-1}} \Delta_{I_{s''}^{(k-1)}} \Big). \qquad (3.16)
\end{aligned}
$$

Analogous rewriting is performed in the case the defect stays close to the right boundary of $\tilde{I}^{(k)}_i$ and $\tilde{I}^{(k)}_{i+1}$ has a defected $(k-1)$–block within distance $3l_k^{\alpha'/\alpha}$ from its left boundary.

iii) No modification case. If during the adjustment procedure the segment $\tilde{I}^{(k)}_i$ remains unchanged, then the corresponding term in (3.12) is not modified.

As long as the adjustment procedure calls on the rearrangement of blocks we continue rewriting the factors as described above, taking into account all modifications which have been performed up to this current step. After a finite number of steps the procedure is stopped and the k–blocks are renamed from left to right as $I^{(k)}_r$.

When all modifications are completed we can write the r.h.s. of the last equality in (3.12) as follows:

$$
\sum_{(I^{(k)},\{D^{(k)}_r\}_r)} \prod_r c(I^{(k)}_r) \prod_r \Big(\prod_{s: I_s^{(k-1)}\subset I^{(k)}_r} \gamma_{I_s^{(k-1)}} + \widehat{\zeta}(I^{(k)}_r) \prod_{\substack{s: I_s^{(k-1)}\subset I^{(k)}_r \\ s\notin D^{(k)}_r}} \gamma_{I_s^{(k-1)}} \prod_{s'\in D^{(k)}_r} \Delta_{I_{s'}^{(k-1)}} \Big), \qquad (3.17)
$$

where, for the fixed sequence $I^{(k-1)}$ (see just before (3.12)), we have:

- the sum $\sum_{(I^k,\{D^{(k)}_r\}_r)}$ is taken over all possible choices of pairs $(I^{(k)},\{D^{(k)}_r\}_r)$ such that

1) $I^{(k)} \equiv I^{(k)}(\{D^{(k)}_r\}_i) = \{I^{(k)}_r\}_r$ is admissible from $I^{(k-1)}$. It depends on the location of the defected $(k-1)$–blocks. The set of indices of defected $(k-1)$–blocks in $I^{(k-1)}$ equals to $\cup_r D^{(k)}_r$;

2) $D^{(k)}_r$ is the set of indices of the defected $(k-1)$–blocks within $I^{(k)}_r$, for each r (we use the same notation for the $\{D^{(k)}_r\}$ before and after the adjustment procedure, but some of these sets might have merged; of course $\cup_i D^{(k)}_i$ does not change during the adjustment procedure;

- the constants $c(I^{(k)}_r) \geq 0$ depend on the all itinerary $J^{(k)}$, and

$$
\sum_{(I^k,\{D^{(k)}_r\}_r)} \prod_r c(I^{(k)}_r) = 1;
$$

- If $k \geq 2$, $\widehat{\zeta}(I_r^{(k)})$ satisfies the following bound:

$$\widehat{\zeta}(I_r^{(k)}) \leq c_2^{-4} c_{8\lfloor l_{k-1}^{\alpha-1} \rfloor}^{-|D_r^{(k)}|} \leq \frac{16}{(\ln 2)^4} \left(\frac{8\lfloor l_{k-1}^{\alpha-1} \rfloor}{\ln 2} \right)^{|D_r|}, \tag{3.18}$$

while for $k = 1$ we should replace $\lfloor l_{k-1}^{\alpha-1} \rfloor$ by l_1 on the r.h.s. of (3.18). To show (3.18) we consider the case when three adjacent intervals merge in the adjustment procedure. The other cases are similar and are left to the reader. Assuming that three blocks $\tilde{I}_i^{(k)}, \tilde{I}_{i+1}^{(k)}$ and $\tilde{I}_{i+2}^{(k)}$ merge, and applying repeatedly (3.16) first to the terms corresponding to $\tilde{I}_i^{(k)}$ and $\tilde{I}_{i+1}^{(k)}$, and then to $\tilde{I}_{i+2}^{(k)}$, we get that the largest corresponding constant is

$$c_2^{-4} c_{j_{i+1}^k - \hat{\jmath}_i^k - 1}^{-|D_i^{(k)}|} c_{\hat{\jmath}_{i+2}^k - \hat{\jmath}_{i+1}^k - 1}^{-|D_{i+1}^{(k)}|} c_{\hat{\jmath}_{i+3}^k - \hat{\jmath}_{i+2}^k - 1}^{-|D_{i+3}^{(k)}|}. \tag{3.19}$$

On the other hand, the estimate (2.8) gives the uniform bound:

$$\hat{\jmath}_{i+1}^k - \hat{\jmath}_i^k - 1 \leq \frac{3l_k + 6l_{k-1}}{l_{k-1} - (2\lfloor l_{k-1}^{\alpha'/\alpha} \rfloor + 6l_{k-2})} \leq 8\lfloor l_{k-1}^{\alpha-1} \rfloor, \tag{3.20}$$

for $k \geq 2$, provided l_1 is large. For $k = 1$ we have $\hat{\jmath}_{i+1}^1 - \hat{\jmath}_i^1 - 1 \leq 3l_1 + 6 \leq 8l_1$. Recalling that $c_N^{-1} \leq N/\ln 2$, (3.20) and (3.19) imply (3.18).

For $I_i^{(k)}$ such that $|D_i^{(k)}| = 1$, say $D_i^{(k)} = \{z\}$, we define

$$g_{I_i^{(k)}} = \prod_{e \in \mathbb{E}^*(I_i^{(k)})} p_e^{\omega_e} q_e^{1-\omega_e} \mathbb{I}_{\omega \in G_z^+(I_i^{(k)})}, \qquad d_{I_i^{(k)}} = \prod_{e \in \mathbb{E}^*(I_i^{(k)})} p_e^{\omega_e} q_e^{1-\omega_e} \mathbb{I}_{\omega \in B_z^+(I_i^{(k)})},$$

where $\mathbb{E}^*(I_i^{(k)}) = \mathbb{E}(I_i^{(k)}) \backslash \cup_s \mathbb{E}(I_s^{(k-1)})$. When $|D_i^{(k)}| > 1$ we define $g_{I_i^{(k)}} \equiv 0$, $d_{I_i^{(k)}} = \prod_{e \in \mathbb{E}^*(I_i^{(k)})} p_e^{\omega_e} q_e^{1-\omega_e}$.

Thus, for $I^{(k-1)}$ fixed and $I^{(k)} = \{I_r^{(k)}\}_r$, obtained from $I^{(k-1)}$, with the collection of sets of defected indices $D \equiv D^{(k)} = \{D_r^{(k)}\}_r$, we write

$$\prod_{I_r^{(k)}} \Big(\prod_{s: I_s^{(k-1)} \subset I_r^{(k)}} \gamma_{I_s^{(k-1)}} + \widehat{\zeta}(I_r^{(k)}) \prod_{\substack{s: I_s^{(k-1)} \subset I_r^{(k)} \\ s \notin D_r^{(k)}}} \gamma_{I_s^{(k-1)}} \prod_{s' \in D_r^{(k)}} \Delta_{I_{s'}^{(k-1)}} \Big) \nu^o_{J_L^{(k-1)}}$$

$$\prod_{I_r^{(k)}} \big(\prod_{s: I_s^{(k-1)} \subset I_r^{(k)}} \gamma_{I_s^{(k-1)}} + \widehat{\zeta}(I_r^{(k)}) \prod_{\substack{s: I_s^{(k-1)} \subset I_r^{(k)} \\ s \notin D_r^{(k)}}} \gamma_{I_s^{(k-1)}} \prod_{s' \in D_r^{(k)}} \Delta_{I_{s'}^{(k-1)}} \big) (g_{I_r^{(k)}} + d_{I_r^{(k)}}) \, \nu^o_{J_L^{(k)}}$$

$$= \prod_{I_r^{(k)}} \Big(\gamma_{I_r^{(k)}} + \Delta_{I_r^{(k)}} \Big) \nu^o_{J_L^{(k)}}, \tag{3.21}$$

where (products over s carry the condition $I_s^{(k-1)} \subset I_r^{(k)}$)

$$\begin{aligned} \gamma_{I_r^{(k)}} &= (g_{I_r^{(k)}} + d_{I_r^{(k)}}) \prod_s \gamma_{I_s^{(k-1)}} + \widehat{\zeta}(I_r^{(k)})\, g_{I_r^{(k)}} \prod_{s \notin D_r^{(k)}} \gamma_{I_s^{(k-1)}} \prod_{s' \in D_r^{(k)}} \Delta_{I_{s'}^{(k-1)}} & (3.22) \\ \Delta_{I_r^{(k)}} &= \widehat{\zeta}(I_r^{(k)})\, d_{I_r^{(k)}} \prod_{s \notin D_r^{(k)}} \gamma_{I_s^{(k-1)}} \prod_{s' \in D_r^{(k)}} \Delta_{I_{s'}^{(k-1)}}. & (3.23) \end{aligned}$$

and $\nu^o_{J_L^{(k)}}$ is the marginal of $\nu^o_{J_L^{(k-1)}}$ (or of ν_L) with all variables corresponding to the edges $e \in \cup_i \mathbb{E}(I_i^{(k)})$ integrated out.

Thus, taking (3.21) into account, (3.9) can be continued as:

$$\begin{aligned} \nu_L &= \sum_{I^{(k-1)} \in \mathcal{I}^{(k-1)}} c(I^{k-1}) \sum_{\{D_r\}_r} \prod_{I_r^{(k)}} c(I_r^{(k)}) \prod_{I_r^{(k)}} \Big(\gamma_{I_r^{(k)}} + \Delta_{I_r^{(k)}} \Big) \nu_{J_L^{(k)}} \\ &= \sum_{I^{(k-1)} \in \mathcal{I}^{(k)}} c(I^k) \prod_{I_r^{(k)} \in \mathcal{I}^{(k)}} \Big(\gamma_{I_r^{(k)}} + \Delta_{I_r^{(k)}} \Big) \nu^o_{J_L^{(k)}}. & (3.24) \end{aligned}$$

Estimates. First of all we observe that using (3.18) and (3.23) we get:

$$\Delta_{I_r^{(k)}}(\omega) \leq \frac{16}{(\ln 2)^4} \left(\frac{8 \lfloor l_{k-1}^{\alpha-1} \rfloor}{\ln 2} \right)^{|D_r^{(k)}|} d_{I_r^{(k)}}(\omega) \prod_{s \notin D_r^{(k)}} \gamma_{I_s^{(k-1)}}(\omega) \prod_{s' \in D_r^{(k)}} \Delta_{I_{s'}^{(k-1)}}(\omega), \qquad (3.25)$$

for $k \geq 2$, while in the case $k = 1$, $\lfloor l_{k-1}^{\alpha-1} \rfloor$ is replaced by l_1.

Given an itinerary J we will define the following sets:

$$\begin{aligned} \widetilde{D}_{r,0}^{(k)}(J) &\equiv \widetilde{D}_{r,0}^{(k)} = \{s \colon I_s^{(k-1)} \subset I_r^{(k)}, \text{all nearest neighbour edges in } I_s^{(k-1)} \text{ are open}\}, \\ \widetilde{D}_{r,1}^{(k)}(J) &\equiv \widetilde{D}_{r,1}^{(k)} = \{s \notin D_r^{(k)} \colon I_s^{(k-1)} \subset I_r^{(k)}, \text{some nearest neighbour edges in } I_s^{(k-1)} \text{ are closed}\}. \end{aligned}$$

We will use the following inequality:

$$\kappa^{\mathcal{C}_{I_r^{(k)}}} \leq \prod_{\substack{s: I_s^{(k-1)} \subset I_r^{(k)} \\ s \in \widetilde{D}_{r,1}^{(k)}}} \kappa^{\mathcal{C}_{I_s^{(k-1)}}} \prod_{\substack{s': I_{s'}^{(k-1)} \subset I_r^{(k)} \\ s' \in D_r^{(k)}}} \kappa^{\mathcal{C}_{I_{s'}^{(k-1)}}}, \qquad (3.26)$$

where it is taken into account that presence of blocks $I_s^{(k-1)} \subset I_r^{(k)}$, $s \in \widetilde{D}_{r,0}^{(k)}(J)$ does not increase the total number of disjoint connected clusters.

Using (2.10), (3.3), (3.4) and (3.25), we get, uniformly in ω:

$$\begin{aligned}
&\kappa^{\mathcal{C}_{I_r^{(k)}}} \Delta_{I_r^{(k)}} \\
\leq\ & \kappa^{\mathcal{C}_{I_r^{(k)}}} \frac{16}{(\ln 2)^4} \left(\frac{8 \left\lfloor l_{k-1}^{\alpha-1} \right\rfloor}{\ln 2} \right)^{|D_r^{(k)}|} d_{I_r^{(k)}} \prod_{\substack{s: I_s^{(k-1)} \subset I_r^{(k)} \\ s \notin D_r^{(k)}}} \gamma_{I_s^{(k-1)}} \prod_{\substack{s': I_{s'}^{(k-1)} \subset I_r^{(k)} \\ s' \in D_r^{(k)}}} \Delta_{I_{s'}^{(k-1)}} \\
\leq\ & \kappa^{\mathcal{C}_{I_r^{(k)}}} \frac{16}{(\ln 2)^4} \left(\frac{8 \left\lfloor l_{k-1}^{\alpha-1} \right\rfloor}{\ln 2} \right)^{|D_r^{(k)}|} d_{I_r^{(k)}} \prod_{s \in \widetilde{D}_{r,0}^{(k)}(J)} \gamma_{I_{s''}^{(k-1)}} \prod_{s \in \widetilde{D}_{r,1}^{(k)}(J)} \gamma_{I_s^{(k-1)}} \prod_{s' \in D_r^{(k)}} \Delta_{I_{s'}^{(k-1)}} \\
\leq\ & \frac{16}{(\ln 2)^4} \left(\frac{8 \left\lfloor l_{k-1}^{\alpha-1} \right\rfloor}{\ln 2} \right)^{|D_r^{(k)}|} d_{I_r^{(k)}} \prod_{\substack{s: I_s^{(k-1)} \subset I_r^{(k)} \\ s \in \widetilde{D}_{r,1}^{(k)}(J)}} \kappa^{\mathcal{C}_{I_s^{(k-1)}}} \gamma_{I_s^{(k-1)}} \prod_{\substack{s': I_{s'}^{(k-1)} \subset I_r^{(k)} \\ s' \in D_r^{(k)}}} \kappa^{\mathcal{C}_{I_{s'}^{(k-1)}}} \Delta_{I_{s'}^{(k-1)}}. \qquad (3.27)
\end{aligned}$$

Recall that, by construction, the variables $d_{I_r^{(k)}}$, $\gamma_{I_s^{(k-1)}}$, $\Delta_{I_{s'}^{(k-1)}}$, with $s \neq s'$, all depend on disjoint sets of edges. For the last inequality of (3.27) we used (3.26) and the trivial bound $\prod_{s \in \widetilde{D}_{r,0}^{(k-1)}(J)} \gamma_{I_{s''}^{(k-1)}}(\omega) \leq 1$. Having (3.27), we consider two cases $|D_r^{(k)}| = 1$ and $|D_r^{(k)}| > 1$. If $|D_r^{(k)}| > 1$ we use the trivial bound $d_{I_r^{(k)}} \leq 1$ and properties stated in 1)-3) at scale $(k-1)$: $\kappa^{\mathcal{C}_{I_s^{(k-1)}}} \gamma_{I_s^{(k-1)}} \leq 1$ for $s \in \widetilde{D}_{r,1}^{(k)}(J)$, and $\kappa^{\mathcal{C}_{I_{s'}^{(k-1)}}} \Delta_{I_{s'}^{(k-1)}} \leq l_{k-1}^{-\delta}$ for $s' \in D_r^{(k)}$, and we get that the r.h.s. of last inequality of (3.27) is bounded by

$$\frac{128}{(\ln 2)^4} l_{k-1}^{(\alpha-1-\delta)|D_r^{(k)}|} \leq \frac{128}{(\ln 2)^4} l_{k-1}^{2(\alpha-1-\delta)} \leq l_k^{-\delta}.$$

If $|D_r^{(k)}| = 1$ we again use $\kappa^{\mathcal{C}_{I_s^{(k-1)}}} \gamma_{I_s^{(k-1)}} \leq 1$ and $\kappa^{\mathcal{C}_{I_{s'}^{(k-1)}}} \Delta_{I_{s'}^{(k-1)}} \leq l_{k-1}^{-\delta}$ as previously, but now we use Lemma 2.1 to estimate $d_{I_r^{(k)}}$. In this case the r.h.s. of last inequality of (3.27) is bounded by

$$\frac{128}{(\ln 2)^5} l_{k-1}^{-\delta+\alpha-1} l_{k-1}^{-\beta(1-\eta)(\alpha'-1)} \leq l_k^{-\delta},$$

and we get the estimate (3.3).

Now we estimate $\kappa^{\mathcal{C}_{I_r^{(k)}}(\omega)} \gamma_{I_r^{(k)}}(\omega)$. First observe that for each ω only one of two summands in (3.22)

$$\left(g_{I_r^{(k)}} + d_{I_r^{(k)}}\right) \prod_{s: I_s^{(k-1)} \subset I_r^{(k)}} \gamma_{I_s^{(k-1)}} \quad \text{or} \quad \widehat{\zeta}(I_r^{(k)})\, g_{I_r^{(k)}} \Delta_{I_{s'}^{(k-1)}} \prod_{\substack{s: I_s^{(k-1)} \subset I_r^{(k)} \\ s \notin D_r}} \gamma_{I_s^{(k-1)}} \qquad (3.28)$$

is non-zero, corresponding to the cases $D_r^{(k)} = \emptyset$ or $D_r^{(k)} = \{s'\}$. The first term can be estimated (recall that we are in the case $\widetilde{D}_{r,1}^{(k)}(J) \neq \emptyset$):

$$\begin{aligned}
\kappa^{\mathcal{C}_{I_r^{(k)}}}\big(g_{I_r^{(k)}} + d_{I_r^{(k)}}\big) \prod_{s: I_s^{(k-1)} \subset I_r^{(k)}} \gamma_{I_s^{(k-1)}} &= \kappa^{\mathcal{C}_{I_r^{(k)}}}\big(g_{I_r^{(k)}} + d_{I_r^{(k)}}\big) \prod_{s'' \in \widetilde{D}_{r,0}^{(k)}(J)} \gamma_{I_{s''}^{(k-1)}} \prod_{s \in \widetilde{D}_{r,1}^{(k)}(J)} \gamma_{I_s^{(k-1)}} \\
&\leq \prod_{s \in \widetilde{D}_{r,1}^{(k)}(J)} \kappa^{\mathcal{C}_{I_s^{(k-1)}}} \gamma_{I_s^{(k-1)}} \leq 1,
\end{aligned} \tag{3.29}$$

where we used trivial bound $\gamma_{I_{s''}^{(k-1)}}(\omega) \leq p_1^{|I_{s''}^{(k-1)}|} < 1$ for $s'' \in \widetilde{D}_{r,0}^{(k)}$. For the second term we have:

$$\begin{aligned}
& \kappa^{\mathcal{C}_{I_r^{(k)}}} \widehat{\zeta}(I_r^{(k)})\, g_{I_r^{(k)}} \Delta_{I_{s'}^{(k-1)}} \prod_{\substack{s: I_s^{(k-1)} \subset I_r^{(k)} \\ s \notin D_r^{(k)}}} \gamma_{I_s^{(k-1)}} \\
= \;& \kappa^{\mathcal{C}_{I_r^{(k)}}} \widehat{\zeta}(I_r^{(k)})\, g_{I_r^{(k)}} \Delta_{I_{s'}^{(k-1)}} \prod_{s \in \widetilde{D}_{r,0}^{(k)}(J)} \gamma_{I_{s''}^{(k-1)}} \prod_{s \in \widetilde{D}_{r,1}^{(k)}(J)} \gamma_{I_s^{(k-1)}} \\
\leq \;& \kappa^{\mathcal{C}_{I_{s'}^{(k-1)}}} \Delta_{I_{s'}^{(k-1)}} \prod_{s \in \widetilde{D}_{r,1}^{(k)}(J)} \kappa^{\mathcal{C}_{I_s^{(k-1)}}} \gamma_{I_s^{(k-1)}} \leq 1.
\end{aligned} \tag{3.30}$$

From (3.29) and (3.30) we get (3.4). To conclude the proof when $k = 1$, we recall the modification of (3.25) in this case, which, together with (3.27) for $k = 1$, and the estimate (3.8), gives

$$\kappa^{\mathcal{C}_{I_r^{(1)}}} \Delta_{I_r^{(1)}}(\omega) \leq \frac{16}{(\ln 2)^4} \left(\frac{8 (\ln 2)^4 l_1^{-\delta}}{128} \right)^{|D_r^{(1)}|} \leq l_1^{-\delta}$$

for every $|D_r^{(1)}| \geq 1$. The other estimates (3.29) and (3.30) follow exactly in the same way. This concludes the proof of Proposition 3.1.□

Since $L = l_M$, at the M–th step of any itinerary, there will be only two intervals $I_1^{(M)}, I_2^{(M)}$. From (3.3) with $k = M$, we easily see that

$$\mu_{\kappa,L}^f \Big(x \rightsquigarrow y, \text{ for some } x \in \big[-L, -L + \lfloor L^{\alpha'/\alpha} \rfloor\big] \text{ and some } y \in \big[L - \lfloor L^{\alpha'/\alpha} \rfloor, L\big] \Big) \geq 1 - 2L^{-\delta}. \tag{3.31}$$

4 Cluster of the origin

The occurrence of the event in the l.h.s. of (3.31) does not imply that the event

$$0 \rightsquigarrow y, \text{ for some } y \in \big[L - \lfloor l_M^{\alpha'/\alpha} \rfloor, L\big]$$

also occurs. Next we give the uniform lower bound for the measure of this last event.

Recalling that $j_z^k = \min\{j : z \in I_j^{(k-1)}\}$, for any $1 \le k \le M$ we define the following events:

$$\psi^{(k)} = \bigcap_{i=j_0^k - \lfloor \lfloor l_k^{\alpha'/\alpha} \rfloor / l_{k-1} \rfloor}^{j_0^k + \lfloor \lfloor l_k^{\alpha'/\alpha} \rfloor / l_{k-1} \rfloor} G(I_i^{(k-1)}) \tag{4.1}$$

and consider

$$\Psi_L = \bigcap_{j=1}^{M} \psi^{(j)}. \tag{4.2}$$

Notice that we have at most two M–blocks, denoted by $I_i^{(M)}$, where $1 \le i \le s$ and $s(\omega) \in \{1, 2\}$.

The occurrence of $\bigcap_{k=1}^{n} \psi^{(k)}$, $1 \le n \le M$ implies that the origin 0 is the right (resp. left) end–vertex of a $(n-1)$–block $I_{j_0^n}^{(n-1)}$ (resp. $I_{j_0^n+1}^{(n-1)}$) for each n, since no adjustments are performed in this case, and necessarily it belongs to the pedestals $\Upsilon(I_{j_0^n}^{(n-1)})$ and $\Upsilon(I_{j_0^n+1}^{(n-1)})$ for any $1 \le n \le M$. In particular, for $\omega \in \Psi_L$ we have $s(\omega) = 2$. Moreover, in the event $\Psi_L \cap G(I_1^{(M)}) \cap G(I_2^{(M)})$, the origin 0 belongs to the open cluster connecting vertices x and y as described above.

Finally, using (3.3) and FKG, and under our choice of parameters $1 < \alpha' < \alpha < 2$, p, l_1 we get

$$\mu_{\kappa,L}^{w}(\Psi_L \cap G(I_1^{(M)}) \cap G(I_2^{(M)})) \ge p^{\lfloor l_1^{\alpha'/\alpha} \rfloor} \prod_{k=2}^{M} (1 - l_{k-1}^{-\delta})^{\lfloor \lfloor l_k^{\alpha'/\alpha} \rfloor / l_{k-1} \rfloor} > c(\alpha, \alpha', p, l_1) > 0$$

since, by (2.19) we have

$$\sum_{k=2}^{\infty} l_{k-1}^{-\delta} \lfloor \lfloor l_k^{\alpha'/\alpha} \rfloor / l_{k-1} \rfloor \le \sum_{k=2}^{\infty} l_{k-1}^{-\delta+\alpha'-1} < \infty \ .$$

This completes the proof of the Theorem 1.1. □

References

[AN] M. Aizenman and C. M. Newman. Discontinuity of the percolation density in one-dimensional $1/|x-y|^2$ models. Commun. Math. Phys. **107**, 611 (1986).

[ACCN] M. Aizenman, J. T. Chayes, L. Chayes and C. M. Newman. Discontinuity of the magnetization in one-dimensional $1/|x-y|^2$ Ising and Potts models. Journ. Stat. Phys. **50**, 1-40 (1988).

[BHS] N. Berger, C. Hoffman, V. Sidoravicius, Non-uniqueness for specifications in $\ell^{2+\epsilon}$. Preprint PR/0312344 v4 (2005)

[CMR] M. Cassandro, I. Merola, U. Rozikov. Phase cohexistence in one dimensional Ising models with long range interactions. (In preparation)

[F] C. M. Fortuin. On the random-cluster model: II. The percolation model. Physica **58**, 393 (1972).

[FK] C. M. Fortuin and P. W. Kasteleyn. On the random-cluster model: I. Introduction and relation to other models. Physica **57**, 536 (1972).

[FS] J. Fröhlich and T. Spencer. The Kosterlitz - Thouless transition in two-dimensional Abelian spin system and the Coulomb gas. Commun. Math. Phys. **81**, 527 (1981).

[FS1] J. Fröhlich and T. Spencer. The phase transition $1/r^2$. Commun. Math. Phys. **84**, 87 (1982).

[G1] G. Grimmett. The stochastic random-cluster process and the uniqueness of random-cluster measures. Ann. Prob. **23**, 1461 (1995).

[G2] G. Grimmett. *The Random-Cluster Model.* Springer. (2006)

[IN] J. Z. Imbrie and C. M. Newman. An Intermediate Phase with Slow Decay of Correlations in One Dimension $1/\left|x-y\right|^2$ Percolation, Ising and Potts Models. Commum. Math. Phys. **118**, 303 (1988).

[JO] A. Johansson and A. Öberg. Square summability of variations of g–functions and uniqueness in g–measures. Math. Res. Lett. **10**, No 5-6, 587-601 (2003).

[M] Domingos H. U. Marchetti. Upper bound on the truncated connectivity in one-dimensional $\beta/\left|x-y\right|^2$ percolation models at $\beta > 1$. Rev. Math. Phys. **7**, 723-742 (1995).

[KMP] A. Klein, D. H. U. Marchetti and J. F. Perez. Power-law fall off in the Kosterlitz–Thouless phase of a two-dimensional lattice Coulomb gas. Journ. Stat. Phys. **60**, 137-166 (1990).

[NS] C. M. Newman and L. S. Schulman. One dimensional $1/|x-y|^s$ percolation models: the existence of a transition for $s \leq 2$, Commun. Math. Phys. **104**, 547 (1986).